

Constructing (ω_1, β) -morasses for $\omega_1 \leq \beta$

Bernhard Irrgang

December 16, 2010

Abstract

Let $\kappa \in \text{Card}$ and $L_\kappa[X]$ be such that the fine structure theory, condensation and $\text{Card}^{L_\kappa[X]} = \text{Card} \cap \kappa$ hold. Then it is possible to prove the existence of morasses. In particular, I will prove that there is a κ -standard morass, a notion that I introduced in a previous paper. This shows the consistency of (ω_1, β) -morasses for all $\beta \geq \omega_1$.

1 Introduction

R. Jensen formulated in the 1970's the concept of an (ω_α, β) -morass whereby objects of size $\omega_{\alpha+\beta}$ could be constructed by a directed system of objects of size less than ω_α . He defined the notion of an (ω_α, β) -morass only for the case that $\beta < \omega_\alpha$. I introduced in a previous paper [Irr2] a definition of an (ω_1, β) -morass for the case that $\omega_1 \leq \beta$.

This definition of an (ω_1, β) -morass for the case that $\omega_1 \leq \beta$ seems to be an axiomatic description of the condensation property of Gödel's constructible universe L and the whole fine structure theory of it. I was, however, not able to formulate and prove this fact in form of a mathematical statement. Therefore, I defined a seemingly innocent strengthening of the notion of an (ω_1, β) -morass, which I actually expect to be equivalent to the notion of (ω_1, β) -morass. I call this strengthening an $\omega_{1+\beta}$ -standard morass. As will be seen, if we construct a morass in the usual way in L , the properties of a standard morass hold automatically.

Using the notion of a standard morass, I was able to prove a theorem which can be interpreted as saying that standard morasses fully cover the condensation property and fine structure of L . More precisely, I was able to show the following [Irr2]

Theorem

Let $\kappa \geq \omega_1$ be a cardinal and assume that a κ -standard morass exists. Then there exists a predicate X such that $\text{Card} \cap \kappa = \text{Card}^{L_\kappa[X]}$ and $L_\kappa[X]$ satisfies amenability, coherence and condensation.

Let me explain this. The predicate X is a sequence $X = \langle X_\nu \mid \nu \in S^X \rangle$ where $S^X \subseteq \text{Lim} \cap \kappa$, and $L_\kappa[X]$ is endowed with the following hierarchy: Let $I_\nu = \langle J_\nu^X, X \upharpoonright \nu \rangle$ for $\nu \in \text{Lim} - S^X$ and $I_\nu = \langle J_\nu^X, X \upharpoonright \nu, X_\nu \rangle$ for $\nu \in S^X$ where $X_\nu \subseteq J_\nu^X$ and

$$J_0^X = \emptyset$$

$$J_{\nu+\omega}^X = rud(I_\nu^X)$$

$$J_\lambda^X = \bigcup \{J_\nu^X \mid \nu \in \lambda\} \text{ for } \lambda \in Lim^2 := Lim(Lim),$$

where $rud(I_\nu^X)$ is the rudimentary closure of $J_\nu^X \cup \{J_\nu^X\}$ relative to $X \upharpoonright \nu$ if $\nu \in Lim - S^X$ and relative to $X \upharpoonright \nu$ and X_ν if $\nu \in S^X$. Now, the properties of $L_\kappa[X]$ are defined as follows:

(Amenability) The structures I_ν are amenable.

(Coherence) If $\nu \in S^X$, $H \prec_1 I_\nu$ and $\lambda = sup(H \cap On)$, then $\lambda \in S^X$ and $X_\lambda = X_\nu \cap J_\lambda^X$.

(Condensation) If $\nu \in S^X$ and $H \prec_1 I_\nu$, then there is some $\mu \in S^X$ such that $H \cong I_\mu$.

Moreover, if we let $\beta(\nu)$ be the least β such that $J_{\beta+\omega}^X \models \nu$ singular, then $S^X = \{\beta(\nu) \mid \nu \text{ singular in } I_\kappa\}$.

As will be seen, these properties suffice to develop the fine structure theory. In this sense, the theorem shows indeed what I claimed. In the present paper, I shall show the converse:

Theorem

If $L_\kappa[X]$, $\kappa \in Card$, satisfies condensation, coherence, amenability, $S^X = \{\beta(\nu) \mid \nu \text{ singular in } I_\kappa\}$ and $Card^{L_\kappa[X]} = Card \cap \kappa$, then there is a κ -standard morass.

Since L itself satisfies the properties of $L_\kappa[X]$, this also shows that the existence of κ -standard morasses and (ω_1, β) -morasses is consistent for all $\kappa \geq \omega_2$ and all $\beta \geq \omega_1$.

Most results that can be proved in L from condensation and the fine structure theory also hold in the structures $L_\kappa[X]$ of the above form. As examples, I proved in my dissertation the following two theorems whose proofs can also be seen as applications of morasses:

Theorem

Let $\lambda \geq \omega_1$ be a cardinal, $S^X \subseteq Lim \cap \lambda$, $Card \cap \lambda = Card^{L_\lambda[X]}$ and $X = \langle X_\nu \mid \nu \in S^X \rangle$ be a sequence such that amenability, coherence, condensation and $S^X = \{\beta(\nu) \mid \nu \text{ singular in } I_\kappa\}$ hold. Then \square_κ holds for all infinite cardinals $\kappa < \lambda$.

Theorem

Let $S^X \subseteq Lim$ and $X = \langle X_\nu \mid \nu \in S^X \rangle$ be a sequence such that amenability, coherence, condensation and $S^X = \{\beta(\nu) \mid \nu \text{ singular in } L[X]\}$ hold. Then the weak covering lemma holds for $L[X]$. That is, if there is no non-trivial, elementary embedding $\pi : L[X] \rightarrow L[X]$, $\kappa \in Card^{L[X]} - \omega_2$ and $\tau = (\kappa^+)^{L[X]}$, then

$$\tau < \kappa^+ \quad \Rightarrow \quad cf(\tau) = card(\kappa).$$

The present paper is a part of my dissertation [Irr1]. I thank Dieter Donder for being my adviser, Hugh Woodin for an invitation to Berkeley, where part of the work was done, and the DFG-Graduiertenkolleg ‘‘Sprache, Information, Logik’’ in Munich for their support.

2 The inner model $L[X]$

We say a function $f : V^n \rightarrow V$ is rudimentary for some structure $\mathfrak{W} = \langle W, X_i \rangle$ if it is generated by the following schemata:

$$\begin{aligned}
 f(x_1, \dots, x_n) &= x_i \text{ for } 1 \leq i \leq n \\
 f(x_1, \dots, x_n) &= \{x_i, x_j\} \text{ for } 1 \leq i, j \leq n \\
 f(x_1, \dots, x_n) &= x_i - x_j \text{ for } 1 \leq i, j \leq n \\
 f(x_1, \dots, x_n) &= h(g_1(x_1, \dots, x_n), \dots, g_n(x_1, \dots, x_n)) \\
 &\text{where } h, g_1, \dots, g_n \text{ are rudimentary} \\
 f(y, x_2, \dots, x_n) &= \bigcup \{g(z, x_2, \dots, x_n) \mid z \in y\} \\
 &\text{where } g \text{ is rudimentary} \\
 f(x_1, \dots, x_n) &= X_i \cap x_j \text{ where } 1 \leq j \leq n.
 \end{aligned}$$

Lemma 1

A function is rudimentary iff it is a composition of the following functions:

$$\begin{aligned}
 F_0(x, y) &= \{x, y\} \\
 F_1(x, y) &= x - y \\
 F_2(x, y) &= x \times y \\
 F_3(x, y) &= \{\langle u, z, v \rangle \mid z \in x \text{ and } \langle u, v \rangle \in y\} \\
 F_4(x, y) &= \{\langle z, u, v \rangle \mid z \in x \text{ and } \langle u, v \rangle \in y\} \\
 F_5(x, y) &= \bigcup x \\
 F_6(x, y) &= \text{dom}(x) \\
 F_7(x, y) &= \in \cap (x \times x) \\
 F_8(x, y) &= \{x[\{z\}] \mid z \in y\} \\
 F_{9+i}(x, y) &= x \cap X_i \text{ for the predicates } X_i \text{ of the structure under consideration.}
 \end{aligned}$$

Proof: See, for example, in [Dev2]. \square

A relation $R \subseteq V^n$ is called rudimentary if there is a rudimentary function $f : V^n \rightarrow V$ such that $R(x_i) \Leftrightarrow f(x_i) \neq \emptyset$.

Lemma 2

Every relation that is Σ_0 over the considered structure is rudimentary.

Proof: Let χ_R be the characteristic function of R . The claim follows from the facts (i)-(vi):

(i) R rudimentary $\Leftrightarrow \chi_R$ rudimentary.

\Leftarrow is clear. Conversely, $\chi_R = \bigcup \{g(y) \mid y \in f(x_i)\}$ where $g(y) = 1$ is constant and $R(x_i) \Leftrightarrow f(x_i) \neq \emptyset$.

(ii) If R is rudimentary, then $\neg R$ is also rudimentary.

Since $\chi_{\neg R} = 1 - \chi_R$.

(iii) $x \in y$ and $x = y$ are rudimentary.

By $x \notin y \Leftrightarrow \{x\} - y \neq \emptyset$, $x \neq y \Leftrightarrow (x - y) \cup (y - x) \neq \emptyset$ and (ii).

(iv) If $R(y, x_i)$ is rudimentary, then $(\exists z \in y)R(z, x_i)$ and $(\forall z \in y)R(z, x_i)$ are rudimentary.

If $R(y, x_i) \Leftrightarrow f(y, x_i) \neq \emptyset$, then $(\exists z \in y)R(z, x_i) \Leftrightarrow \bigcup\{f(z, x_i) \mid z \in y\} \neq \emptyset$. The second claim follows from this by (ii).

(v) If $R_1, R_2 \subseteq V^n$ are rudimentary, then so are $R_1 \vee R_2$ and $R_1 \wedge R_2$.

Because $f(x, y) = x \cup y$ is rudimentary, $(R_1 \vee R_2)(x_i) \Leftrightarrow \chi_{R_1}(x_i) \cup \chi_{R_2}(x_i) \neq \emptyset$ is rudimentary. The second claim follows from that by (ii).

(vi) $x \in X_i$ is rudimentary.

Since $\{x\} \cap X_i \neq \emptyset \Leftrightarrow x \in X_i$. \square

For a converse of this lemma, we define:

A function f is called simple if $R(f(x_i), y_k)$ is Σ_0 for every Σ_0 -relation $R(z, y_k)$.

Lemma 3

A function f is simple iff

- (i) $z \in f(x_i)$ is Σ_0
- (ii) $A(z)$ is $\Sigma_0 \Rightarrow (\exists z \in f(x_i))A(z)$ is Σ_0 .

Proof: If f is simple, then (i) and (ii) hold, because these are instances of the definition. The converse is proved by induction on Σ_0 -formulas. E.g. if $R(z, y_k) : \Leftrightarrow z = y_k$, then $R(f(x_i), y_k) \Leftrightarrow f(x_i) = y_k \Leftrightarrow (\forall z \in f(x_i))(z \in y_k)$ and $(\forall z \in y_k)(z \in f(x_i))$. Thus we need (i) and (ii). The other cases are similar. \square

Lemma 4

Every rudimentary function is Σ_0 in the parameters X_i .

Proof: By induction, one proves that the rudimentary functions that are generated without the schema $f(x_1, \dots, x_n) = X_i \cap x_j$ are simple. For this, one uses lemma 3. But since the function $f(x, y) = x \cap y$ is one of those, the claim holds. \square

Thus every rudimentary relation is Σ_0 in the parameters X_i , but not necessarily Σ_0 with the X_i as predicates. An example is the relation $\{x, y\} \in X_0$.

A structure is said to be rudimentary closed if its underlying set is closed under all rudimentary functions.

Lemma 5

If \mathfrak{M} is rudimentary closed and $H \prec_1 \mathfrak{M}$, then H and the collapse of H are also rudimentary closed.

Proof: That is clear, since the functions F_0, \dots, F_{9+i} are Σ_0 with the predicates X_i . \square

Let T_N be the set of Σ_0 formulae of our language $\{\in, X_1, \dots, X_N\}$ having exactly one free variable. By lemma 2, there is a rudimentary function f for every Σ_0 formula ψ such that $\psi(x_*) \Leftrightarrow f(x_*) \neq \emptyset$. By lemma 1, we have

$$\begin{aligned} x_0 &= f(x_*) = F_{k_1}(x_1, x_2) \\ \text{where } x_1 &= F_{k_2}(x_3, x_4) \\ x_2 &= F_{k_3}(x_5, x_6) \\ \text{and } x_3 &= \dots \end{aligned}$$

Of course, x_* appears at some point.

Therefore, we may define an effective Gödel coding

$$T_N \rightarrow G, \psi_u \mapsto u$$

as follows (m, n possibly $= \star$):

$$\langle k, l, m, n \rangle \in u \Leftrightarrow x_k = F_l(x_m, x_n).$$

Let $\models_{\mathfrak{W}}^{\Sigma_0}(u, x_*) \Leftrightarrow$

ψ_u is a Σ_0 formula with exactly one free variable
and $\mathfrak{W} \models \psi_u(x_*)$.

Lemma 6

If \mathfrak{W} is transitive and rudimentary closed, then $\models_{\mathfrak{W}}^{\Sigma_0}(x, y)$ is Σ_1 -definable over \mathfrak{W} . The definition of $\models_{\mathfrak{W}}^{\Sigma_0}(x, y)$ depends only on the number of predicates of \mathfrak{W} . That is, it is uniform for all structures of the same type.

Proof: Whether $\models_{\mathfrak{W}}^{\Sigma_0}(u, x_*)$ holds, may be computed directly. First, one computes the x_k which only depend on x_* . For those k , $\langle k, l, \star, \star \rangle \in u$. Then one computes the x_i which only depend on x_m and x_n such that $m, n \in \{k \mid \langle k, l, \star, \star \rangle \in u\}$ – etc. Since \mathfrak{W} is rudimentary closed, this process only breaks off, when one has computed $x_0 = f(x_*)$. And $\models_{\mathfrak{W}}^{\Sigma_0}(u, x_*)$ holds iff $x_0 = f(x_*) \neq \emptyset$.

More formally speaking: $\models_{\mathfrak{W}}^{\Sigma_0}(u, x_*)$ holds iff there is some sequence $\langle x_i \mid i \in d \rangle$, $d = \{k \mid \langle k, l, m, n \rangle \in u\}$ such that

$$\langle k, l, m, n \rangle \in u \Rightarrow x_k = F_l(x_m, x_n)$$

and $x_0 \neq \emptyset$.

Hence $\models_{\mathfrak{W}}^{\Sigma_0}$ is Σ_1 . \square

If \mathfrak{W} is a structure, then let $rud(\mathfrak{W})$ be the closure of $W \cup \{W\}$ under the functions which are rudimentary for \mathfrak{W} .

Lemma 7

If \mathfrak{W} is transitive, then so is $rud(\mathfrak{W})$.

Proof: By induction on the definition of the rudimentary functions. \square

Lemma 8

Let \mathfrak{W} be a transitive structure with underlying set W . Then

$$rud(\mathfrak{W}) \cap \mathfrak{P}(W) = Def(\mathfrak{W}).$$

Proof: First, let $A \in Def(\mathfrak{W})$. Then A is Σ_0 over $\langle W \cup \{W\}, X_i \rangle$, i.e. there are parameters $p_i \in W \cup \{W\}$ and some Σ_0 formula φ such that $x \in A \Leftrightarrow \varphi(x, p_i)$. But by lemma 2, every Σ_0 relation is rudimentary. Thus there is a rudimentary function f such that $x \in A \Leftrightarrow f(x, p_i) \neq \emptyset$. Let $g(z, x) = \{x\}$ and define $h(y, x) = \bigcup \{g(z, x) \mid z \in y\}$. Then $h(f(x, p_i), x) = \bigcup \{g(z, x) \mid z \in f(x, p_i)\}$ is rudimentary, $h(f(x, p_i), x) = \emptyset$ if $x \notin A$ and $h(f(x, p_i), x) = \{x\}$ if $x \in A$. Finally, let $H(y, p_i) = \bigcup \{h(f(x, p_i), x) \mid x \in y\}$. Then H is rudimentary and $A = H(W, p_i)$. So we are done.

Conversely, let $A \in \text{rud}(\mathfrak{W}) \cap \mathfrak{P}(W)$. Then there is a rudimentary function f and some $a \in W$ such that $A = f(a, W)$. By lemma 4 and lemma 3, there exists a Σ_0 formula ψ such that $x \in f(a, W) \Leftrightarrow \psi(x, a, W, X_i)$. By Σ_0 absoluteness, $A = \{x \in W \mid W \cup \{W, X_i\} \models \psi(x, a, W, X_i)\}$, since $X_i \subseteq W$. Therefore, there is a formula φ such that $A = \{x \in W \mid \mathfrak{W} \models \varphi(x, a)\}$. \square

Let $\kappa \in \text{Card} - \omega_1$, $S^X \subseteq \text{Lim} \cap \kappa$ and $\langle X_\nu \mid \nu \in S^X \rangle$ be a sequence.

For $\nu \in \text{Lim} - S^X$, let $I_\nu = \langle J_\nu^X, X \upharpoonright \nu \rangle$ and let $I_\nu = \langle J_\nu^X, X \upharpoonright \nu, X_\nu \rangle$ for $\nu \in S^X$ such that

$X_\nu \subseteq J_\nu^X$ where

$$J_0^X = \emptyset$$

$$J_{\nu+\omega}^X = \text{rud}(I_\nu)$$

$$J_\lambda^X = \bigcup \{J_\nu^X \mid \nu \in \lambda\} \text{ if } \lambda \in \text{Lim}^2 := \text{Lim}(\text{Lim}).$$

Obviously, $L_\kappa[X] = \bigcup \{J_\nu^X \mid \nu \in \kappa\}$.

We say that $L_\kappa[X]$ is amenable if I_ν is rudimentary closed for all $\nu \in S^X$.

Lemma 9

- (i) Every J_ν^X is transitive
- (ii) $\mu < \nu \Rightarrow J_\mu^X \in J_\nu^X$
- (iii) $\text{rank}(J_\nu^X) = J_\nu^X \cap \text{On} = \nu$

Proof: That are three easy proofs by induction. \square

Sometimes we need levels between J_ν^X and $J_{\nu+\omega}^X$. To make those transitive, we define

$$G_i(x, y, z) = F_i(x, y) \text{ for } i \leq 8$$

$$G_9(x, y, z) = x \cap X$$

$$G_{10}(x, y, z) = \langle x, y \rangle$$

$$G_{11}(x, y, z) = x[y]$$

$$G_{12}(x, y, z) = \{\langle x, y \rangle\}$$

$$G_{13}(x, y, z) = \langle x, y, z \rangle$$

$$G_{14}(x, y, z) = \{\langle x, y \rangle, z\}.$$

Let

$$S_0 = \emptyset$$

$$S_{\mu+1} = S_\mu \cup \{S_\mu\} \cup \bigcup \{G_i[(S_\mu \cup \{S_\mu\})^3] \mid i \in 15\}$$

$$S_\lambda = \bigcup \{S_\mu \mid \mu \in \lambda\} \text{ if } \lambda \in \text{Lim}.$$

Lemma 10

The sequence $\langle I_\mu \mid \mu \in \text{Lim} \cap \nu \rangle$ is (uniformly) Σ_1 -definable over I_ν .

Proof: By definition $J_\mu^X = S_\mu$ for $\mu \in \text{Lim}$, that is, the sequence $\langle J_\mu^X \mid \mu \in \text{Lim} \cap \nu \rangle$ is the solution of the recursion defining S_μ restricted to Lim . Since the recursion condition is Σ_0 over I_ν , the solution is Σ_1 . It is Σ_1 over I_ν if the existential quantifier can be restricted to J_ν^X . Hence we must prove $\langle S_\mu \mid \mu \in \tau \rangle \in J_\nu^X$ for $\tau \in \nu$. This is done by induction on ν . The base case $\nu = 0$ and the limit step are clear. For the successor step, note that $S_{\mu+1}$ is a rudimentary function of S_μ and μ , and use the rudimentary closedness of J_ν^X .

□

Lemma 11

There are well-orderings $<_\nu$ of the sets J_ν^X such that

- (i) $\mu < \nu \Rightarrow <_\mu \subseteq <_\nu$
- (ii) $<_{\nu+1}$ is an end-extension of $<_\nu$
- (iii) The sequence $\langle <_\mu \mid \mu \in \text{Lim} \cap \nu \rangle$ is (uniformly) Σ_1 -definable over I_ν .
- (iv) $<_\nu$ is (uniformly) Σ_1 -definable over I_ν .
- (v) The function $pr_\nu(x) = \{z \mid z <_\nu x\}$ is (uniformly) Σ_1 -definable over I_ν .

Proof: Define well-orderings $<_\mu$ of S_μ by recursion:

- (I) $<_0 = \emptyset$
- (II) (1) For $x, y \in S_\mu$, let $x <_{\mu+1} y \Leftrightarrow x <_\mu y$
 (2) $x \in S_\mu$ and $y \notin S_\mu \Rightarrow x <_{\mu+1} y$
 $y \in S_\mu$ and $x \notin S_\mu \Rightarrow y <_{\mu+1} x$
 (3) If $x, y \notin S_\mu$, then there is an $i \in 15$ and $x_1, x_2, x_3 \in S_\mu$ such that $x = G_i(x_1, x_2, x_3)$. And there is a $j \in 15$ and $y_1, y_2, y_3 \in S_\mu$ such that $y = G_j(y_1, y_2, y_3)$. First, choose i and j minimal, then x_1 and y_1 , then x_2 and y_2 , and finally x_3 and y_3 .
 Set:
 (a) $x <_{\mu+1} y$ if $i < j$
 $y <_{\mu+1} x$ if $j < i$
 (b) $x <_{\mu+1} y$ if $i = j$ and $x_1 <_\mu y_1$
 $y <_{\mu+1} x$ if $i = j$ and $y_1 <_\mu x_1$
 (c) $x <_{\mu+1} y$ if $i = j$ and $x_1 = y_1$ and $x_2 <_\mu y_2$
 $y <_{\mu+1} x$ if $i = j$ and $x_1 = y_1$ and $y_2 <_\mu x_2$
 (d) $x <_{\mu+1} y$ if $i = j$ and $x_1 = y_1$ and $x_2 = y_2$ and $x_3 <_\mu y_3$
 $y <_{\mu+1} x$ if $i = j$ and $x_1 = y_1$ and $y_2 = x_2$ and $y_3 <_\mu x_3$
- (III) $<_\lambda = \bigcup \{<_\mu \mid \mu \in \lambda\}$

The properties (i) to (v) are obvious. For the Σ_1 -definability, one needs the argument from lemma 10. □

Lemma 12

The rudimentary closed $\langle J_\nu^X, X \upharpoonright \nu, A \rangle$ have a canonical Σ_1 -Skolem function h .

Proof: Let $\langle \psi_i \mid i \in \omega \rangle$ be an effective enumeration of the Σ_0 formulae with three free variables. Intuitively, we would define:

$$h(i, x) \simeq (z)_0$$

for

the $<_\nu$ -least $z \in J_\nu^X$ such that $\langle J_\nu^X, X \upharpoonright \nu, A \rangle \models \psi_i((z)_0, x, (z)_1)$.

Formally, we define:

By lemma 11 (v), let θ be a Σ_0 formula such that

$$w = \{v \mid v <_\nu z\} \Leftrightarrow \langle J_\nu^X, X \upharpoonright \nu, A \rangle \models (\exists t)\theta(w, z, t).$$

Let u_i be the Gödel coding of

$$\theta((s)_1, (s)_0, (s)_2)$$

$$\wedge \psi_i(((s)_0)_0, (s)_3, ((s)_0)_1) \wedge (\forall v \in (s)_1) \neg \psi_i((v)_0, (s)_3, (v)_1)$$

and

$$y = h(i, x) \Leftrightarrow$$

$$(\exists s)((s)_0 = y \wedge (s)_3 = x \wedge \models_{\langle J_\nu^X, X \upharpoonright \nu, A \rangle}^{\Sigma_0} (u_i, s)).$$

This has the desired properties. Note lemma 6! \square

I will denote this Σ_1 -Skolem function by $h_{\nu, A}$. Let $h_\nu := h_{\nu, \emptyset}$.

Let us say that $L_\kappa[X]$ has condensation if the following holds:

If $\nu \in S^X$ and $H <_1 I_\nu$, then there is some $\mu \in S^X$ such that $H \cong I_\mu$.

From now on, suppose that $L_\kappa[X]$ is amenable and has condensation.

Set $I_\nu^0 = \langle J_\nu^X, X \upharpoonright \nu \rangle$ for all $\nu \in Lim \cap \kappa$.

Lemma 13 (Gödel's pairing function)

There is a bijection $\Phi : On^2 \rightarrow On$ such that $\Phi(\alpha, \beta) \geq \alpha, \beta$ for all α, β and $\Phi^{-1} \upharpoonright \alpha$ is uniformly Σ_1 -definable over I_α^0 for all $\alpha \in Lim$.

Proof: Define a well-ordering $<^*$ on On^2 by

$$\langle \alpha, \beta \rangle <^* \langle \gamma, \delta \rangle$$

iff

$$\begin{aligned} &max(\alpha, \beta) < max(\gamma, \delta) \text{ or} \\ &max(\alpha, \beta) = max(\gamma, \delta) \text{ and } \alpha < \gamma \text{ or} \\ &max(\alpha, \beta) = max(\gamma, \delta) \text{ and } \alpha = \gamma \text{ and } \beta < \delta. \end{aligned}$$

Let $\Phi : \langle On^2, <^* \rangle \cong \langle On, < \rangle$. Then Φ may be defined by the recursion

$$\begin{aligned} \Phi(0, \beta) &= sup\{\Phi(\nu, \nu) \mid \nu < \beta\} \\ \Phi(\alpha, \beta) &= \Phi(0, \beta) + \alpha \text{ if } \alpha < \beta \\ \Phi(\alpha, \beta) &= \Phi(0, \alpha) + \alpha + \beta \text{ if } \alpha \geq \beta. \end{aligned}$$

\square

So there is a uniform map from α onto $\alpha \times \alpha$ for all α that are closed under Gödel's pairing function. Such a map exists for all $\alpha \in Lim$. But then we have to give up uniformity.

Lemma 14

For all $\alpha \in Lim$, there exists a function from α onto $\alpha \times \alpha$ that is Σ_1 -definable over I_α^0 .

Proof by induction on $\alpha \in Lim$. If α is closed under Gödel's pairing function, then lemma 13 does the job. Therefore, if $\alpha = \beta + \omega$ for some $\beta \in Lim$, we may assume $\beta \neq 0$. But then there is some over I_α^0 Σ_1 -definable bijection $j : \alpha \rightarrow \beta$. And by the induction hypothesis, there is an over I_β^0 Σ_1 -definable function from β onto $\beta \times \beta$. Thus there exists a Σ_1 formula $\varphi(x, y, p)$ and a parameter $p \in J_\beta^X$ such that there is some $x \in \beta$ satisfying $\varphi(x, y, p)$ for all $y \in \beta \times \beta$. So we

get an over I_β^0 Σ_1 -definable injective function $g : \beta \times \beta \rightarrow \beta$ from the Σ_1 -Skolem function. Hence $f(\langle \nu, \tau \rangle) = g(\langle j(\nu), j(\tau) \rangle)$ defines an injective function $f : \alpha^2 \rightarrow \beta$ which is Σ_1 -definable over I_α^0 . An h which is as needed may be defined by

$$\begin{aligned} h(\nu) &= f^{-1}(\nu) \text{ if } \nu \in \text{rng}(f) \\ h(\nu) &= \langle 0, 0 \rangle \text{ else.} \end{aligned}$$

For $\text{rng}(f) = \text{rng}(g) \in J_\alpha^X$.

Now, assume $\alpha \in \text{Lim}^2$ is not closed under Gödel's pairing function. Then $\nu, \tau \in \alpha$ for $\langle \nu, \tau \rangle = \Phi^{-1}(\alpha)$, and $c := \{z \mid z <^* \langle \nu, \tau \rangle\}$ lies in J_α^X . Thus $\Phi^{-1} \upharpoonright c : c \rightarrow \alpha$ is an over I_α^0 Σ_1 -definable bijection. Pick a $\gamma \in \text{Lim}$ such that $\nu, \tau < \gamma$. Then $\Phi^{-1} \upharpoonright \alpha : \alpha \rightarrow \gamma^2$ is an over I_α^0 Σ_1 -definable injective function. Like in the first case, there exists an injective function $g : \gamma \times \gamma \rightarrow \gamma$ in J_α^X by the induction hypothesis. So $f(\langle \xi, \zeta \rangle) = g(\langle g\Phi^{-1}(\xi), g\Phi^{-1}(\zeta) \rangle)$ defines an over I_α^0 Σ_1 -definable bijection $f : \alpha^2 \rightarrow d$ such that $d := g[g[c] \times g[c]]$. Again, we define h by

$$\begin{aligned} h(\xi) &= f^{-1}(\xi) \text{ if } \xi \in d \\ h(\xi) &= \langle 0, 0 \rangle \text{ else. } \square \end{aligned}$$

Lemma 15

Let $\alpha \in \text{Lim} - \omega + 1$. Then there is some over I_α^0 Σ_1 -definable function from α onto J_α^X . This function is uniformly definable for all α closed under Gödel's pairing function.

Proof: Let $f : \alpha \rightarrow \alpha \times \alpha$ be a surjective function which is Σ_1 -definable over I_α^0 with parameter p . Let p be minimal with respect to the canonical well-ordering such that such an f exists. Define f^0, f^1 by $f(\nu) = \langle f^0(\nu), f^1(\nu) \rangle$ and, by induction, define $f_1 = id \upharpoonright \alpha$ and $f_{n+1}(\nu) = \langle f^0(\nu), f_n \circ f^1(\nu) \rangle$. Let $h := h_\alpha$ be the canonical Σ_1 -Skolem function and $H = h[\omega \times (\alpha \times \{p\})]$. Then H is closed under ordered pairs. For, if $y_1 = h(j_1, \langle \nu_1, p \rangle)$, $y_2 = h(j_2, \langle \nu_2, p \rangle)$ and $\langle \nu_1, \nu_2 \rangle = f(\tau)$, then $\langle y_1, y_2 \rangle$ is Σ_1 -definable over I_α^0 with the parameters τ, p . Hence it is in H . Since H is closed under ordered pairs, we have $H \prec_1 I_\alpha^0$. Let $\sigma : H \rightarrow I_\beta^0$ be the collapse of H . Then $\alpha = \beta$, because $\alpha \subseteq H$ and $\sigma \upharpoonright \alpha = id \upharpoonright \alpha$. Thus $\sigma[f] = f$, and $\sigma[f]$ is Σ_1 -definable over I_α^0 with the parameter $\sigma(p)$. Since σ is a collapse, $\sigma(p) \leq p$. So $\sigma(p) = p$ by the minimality of p . In general, $\pi(h(i, x)) \simeq h(i, \pi(x))$ for Σ_1 -elementary π . Therefore, $\sigma(h(i, \langle \nu, p \rangle)) \simeq h(i, \langle \nu, p \rangle)$ holds in our case for all $i \in \omega$ and $\nu \in \alpha$. But then $\sigma \upharpoonright H = id \upharpoonright H$ and $H = J_\alpha^X$. Thus we may define the needed surjective map by $g \circ f_3$ where

$$\begin{aligned} g(i, \nu, \tau) &= y \text{ if } (\exists z \in S_\tau) \varphi(z, y, i, \langle \nu, p \rangle) \\ g(i, \nu, \tau) &= \emptyset \text{ else.} \end{aligned}$$

Here, S_τ shall be defined as in lemma 10 and $y = h(i, x) \Leftrightarrow (\exists t \in J_\alpha^X) \varphi(t, i, x, y)$. \square

Let $\langle I_\nu^0, A \rangle := \langle J_\nu^X, X \upharpoonright \nu, A \rangle$.

The idea of the fine structure theory is to code Σ_n predicates over large structures in Σ_1 predicates over smaller structures. In the simplest case, one codes the Σ_1 information of the given structure I_β^0 in a rudimentary closed structure $\langle I_\rho^0, A \rangle$. I.e. we want to have something like:

Over I_β^0 , there exists a Σ_1 function f such that

$$f[J_\rho^X] = J_\beta^X.$$

For the Σ_1 formulae φ_i ,

$$\langle i, x \rangle \in A \quad \Leftrightarrow \quad I_\beta^0 \models \varphi_i(f(x))$$

holds. And

$$\langle I_\rho^0, A \rangle \quad \text{is rudimentary closed.}$$

Now, suppose we have such an $\langle I_\rho^0, A \rangle$. Then every $B \subseteq J_\rho^X$ that is Σ_1 -definable over I_β^0 is of the form

$$B = \{x \mid A(i, \langle x, p \rangle)\} \quad \text{for some } i \in \omega, p \in J_\rho^X.$$

So $\langle I_\rho^0, B \rangle$ is rudimentary closed for all $B \in \Sigma_1(I_\beta^0) \cap \mathfrak{P}(J_\rho^X)$.

The ρ is uniquely determined.

Lemma 16

Let $\beta > \omega$ and $\langle I_\beta^0, B \rangle$ be rudimentary closed. Then there is at most one $\rho \in Lim$ such that

$$\langle I_\rho^0, C \rangle \text{ is rudimentary closed for all } C \in \Sigma_1(\langle I_\beta^0, B \rangle) \cap \mathfrak{P}(J_\rho^X)$$

and

$$\text{there is an over } \langle I_\beta^0, B \rangle \text{ } \Sigma_1\text{-definable function } f \text{ such that } f[J_\rho^X] = J_\beta^X.$$

Proof: Assume $\rho < \bar{\rho}$ both had these properties. Let f be an over $\langle I_\beta^0, B \rangle$ Σ_1 -definable function such that $f[J_\rho^X] = J_\beta^X$ and $C = \{x \in J_\rho^X \mid x \notin f(x)\}$. Then $C \subseteq J_\rho^X$ is Σ_1 -definable over $\langle I_\beta^0, B \rangle$. So $\langle I_\rho^0, C \rangle$ is rudimentary closed. But then $C = C \cap J_\rho^X \in J_\rho^X$. Hence there is an $x \in J_\rho^X$ such that $C = f(x)$. From this, the contradiction $x \in f(x) \Leftrightarrow x \in C \Leftrightarrow x \notin f(x)$ follows. \square

The uniquely determined ρ from lemma 16 is called the projectum of $\langle I_\beta^0, B \rangle$.

If there is some over $\langle I_\beta^0, B \rangle$ Σ_1 -definable function f such that $f[J_\rho^X] = J_\beta^X$, then $h_{\beta, B}[\omega \times (J_\rho^X \times \{p\})] = J_\beta^X$ for a $p \in J_\beta^X$. Using the canonical function $h_{\beta, B}$, we can define a canonical A :

Let p be minimal with respect to the canonical well-ordering such that the above property holds. Define

$$A = \{\langle i, x \rangle \mid i \in \omega \quad \text{and} \quad x \in J_\rho^X \quad \text{and} \quad \langle I_\beta^0, B \rangle \models \varphi_i(x, p)\}.$$

We say p is the standard parameter of $\langle I_\beta^0, B \rangle$ and A the standard code of it.

Lemma 17

Let $\beta > 0$ and $\langle I_\beta^0, B \rangle$ be rudimentary closed. Let ρ be the projectum and A the standard code of it. Then for all $m \geq 1$, the following holds:

$$\Sigma_{1+m}(\langle I_\beta^0, B \rangle) \cap \mathfrak{P}(J_\rho^X) = \Sigma_m(\langle I_\rho^0, A \rangle).$$

Proof: First, let $R \in \Sigma_{1+m}(\langle I_\beta^0, B \rangle) \cap \mathfrak{P}(J_\rho^X)$ and let m be even. Let P be a relation being Σ_1 -definable over $\langle I_\beta^0, B \rangle$ with parameter q_1 such that, for $x \in J_\rho^X$, $R(x)$ holds iff $\exists y_0 \forall y_1 \exists y_3 \dots \forall y_{m-1} P(y_i, x)$. Let f be some over $\langle I_\beta^0, B \rangle$ with parameter q_2 Σ_1 -definable function such that $f[J_\rho^X] = J_\beta^X$. Define $Q(z_i, x)$ by $z_i, x \in J_\rho^X$ and $(\exists y_i)(y_i = f(z_i) \text{ and } P(y_i, x))$. Let p be the standard parameter

of $\langle I_\beta^0, B \rangle$. Then, by definition, there is some $u \in J_\rho^X$ such that $\langle q_1, q_2 \rangle$ is Σ_1 -definable in $\langle I_\beta^0, B \rangle$ with the parameters u, p . I.e. there is some $i \in \omega$ such that $Q(z_i, x)$ holds iff $z_i, x \in J_\rho^X$ and $\langle I_\beta^0, B \rangle \models \varphi_i(\langle z_i, x, u \rangle, p)$ – i.e. iff $z_i, x \in J_\rho^X$ and $A(i, \langle z_i, x, u \rangle)$. Analogously there is a $j \in \omega$ and a $v \in J_\rho^X$ such that $z \in \text{dom}(f) \cap J_\rho^X$ iff $z \in J_\rho^X$ and $A(j, \langle z, v \rangle)$. Abbreviate this by $D(z)$. But then, for $x \in J_\rho^X$, $R(x)$ holds iff $\exists y_0 \forall y_1 \exists y_2 \dots \forall y_{m-1} (D(z_0) \wedge D(z_2) \wedge \dots \wedge D(z_{m-2})$ and $(D(z_1) \wedge D(z_3) \wedge \dots \wedge D(z_{m-1}) \Rightarrow Q(z_i, x))$. So the claim holds. If m is odd, then we proceed correspondingly. Thus $\Sigma_{1+m}(\langle I_\beta^0, B \rangle) \cap \mathfrak{P}(J_\rho^X) \subseteq \Sigma_m(\langle I_\rho^0, A \rangle)$ is proved.

Conversely, let φ be a Σ_0 formula and $q \in J_\rho^X$ such that, for all $x \in J_\rho^X$, $R(x)$ holds iff $\langle I_\rho^0, A \rangle \models \varphi(x, q)$. Since $\langle I_\rho^0, A \rangle$ is rudimentary closed, $R(x)$ holds iff $(\exists u \in J_\rho^X)(\exists a \in J_\rho^X)(u \text{ transitive and } x \in u \text{ and } q \in u \text{ and } a = A \cap u \text{ and } \langle u, a \rangle \models \varphi(x, q))$. Write $a = A \cap u$ as formula: $(\forall v \in a)(v \in u \text{ and } v \in A)$ and $(\forall v \in u)(v \in A \Rightarrow v \in a)$. If $m = 1$, we are done provided we can show that this is Σ_2 over $\langle I_\beta^0, B \rangle$. If $m > 1$, the claim follows immediately by induction. The second part is Π_1 . So we only have to prove that the first part is Σ_2 over $\langle I_\beta^0, B \rangle$. By the definition of A , $v \in A$ is Σ_1 -definable over $\langle I_\beta^0, B \rangle$. I.e. there is some Σ_0 formula ψ and some parameter p such that $v \in A \Leftrightarrow \langle I_\beta^0, B \rangle \models (\exists y)\psi(v, y, p)$. Now, we have two cases.

In the first case, there is no over $\langle I_\beta^0, B \rangle$ Σ_1 -definable function from some $\gamma < \rho$ cofinal in β . Then $(\forall v \in a)(v \in A)$ is Σ_2 over $\langle I_\beta^0, B \rangle$, because some kind of replacement axiom holds, and $(\forall v \in a)(\exists y)\psi(v, y, p)$ is over $\langle I_\beta^0, B \rangle$ equivalent to $(\exists z)(\forall v \in a)(\exists y \in z)\psi(v, y, p)$. For $\rho = \omega$, this is obvious. If $\rho \neq \omega$, then $\rho \in \text{Lim}^2$ and we can pick a $\gamma < \rho$ such that $a \in J_\gamma^X$. Let $j : \gamma \rightarrow J_\gamma^X$ an over I_γ Σ_1 -definable surjection, and g an over $\langle I_\beta^0, B \rangle$ Σ_1 -definable function that maps $v \in J_\beta^X$ to $g(v) \in J_\beta^X$ such that $\psi(v, g(v), p)$ if such an element exists. We can find such a function with the help of the Σ_1 -Skolem function. Now, define a function $f : \gamma \rightarrow \beta$ by

$$\begin{aligned} f(\nu) &= \text{the least } \tau < \beta \text{ such that } g \circ j(\nu) \in S_\tau \text{ if } j(\nu) \in a \\ f(\nu) &= 0 \text{ else.} \end{aligned}$$

Since f is Σ_1 , there is, in the given case, a $\delta < \beta$ such that $f[\gamma] \subseteq \delta$. So we have as collecting set $z = S_\delta$, and the equivalence is clear.

Now, let us come to the second case. Let $\gamma < \rho$ be minimal such that there is some over $\langle I_\beta^0, B \rangle$ Σ_1 -definable function g from γ cofinal in β . Then $(\forall v \in a)(\exists y)\psi(v, y, p)$ is equivalent to $(\forall v \in a)(\exists \nu \in \gamma)(\exists y \in S_{g(\nu)})\psi(v, y, p)$. If we define a predicate $C \subseteq J_\rho^X$ by $\langle v, \nu \rangle \in C \Leftrightarrow y \in S_{g(\nu)}$ and $\psi(v, y, p)$, then $\langle I_\beta^0, B \rangle \models (\forall v \in a)(\exists y)\psi(v, y, p)$ is equivalent to $\langle I_\rho^0, C \rangle \models (\forall v \in a)(\exists \nu \in \gamma)(\exists y)((\langle v, \nu \rangle \in C))$. But this holds iff $\langle I_\rho^0, C \rangle \models (\exists w)(w \text{ transitive and } a, \gamma \in w \text{ and } \langle w, C \cap w \rangle \models (\forall v \in a)(\exists \nu \in \gamma)(\exists y)((\langle v, \nu \rangle \in C \cap w))$. Since C is Σ_1 over $\langle I_\beta^0, B \rangle$, $\langle I_\rho^0, C \rangle$ is rudimentary closed by the definition of the projectum. I.e. the statement is equivalent to $\langle I_\rho^0, C \rangle \models (\exists w)(\exists c)(w \text{ transitive and } a, \gamma \in w \text{ and } c = C \cap w \text{ and } \langle w, c \rangle \models (\forall v \in a)(\exists \nu \in \gamma)(\exists y)((\langle v, \nu \rangle \in c))$. So, to prove that this is Σ_2 , it suffices to show that $c = C \cap w$ is Σ_2 . In its full form, this is $(\forall z)(z \in a \Leftrightarrow z \in w \text{ and } z \in C)$. But $z \in C$ is even Δ_1 over $\langle I_\beta^0, B \rangle$ by the definition. So we are finished. \square

Lemma 18

(a) Let $\pi : \langle J_\beta^X, X \upharpoonright \bar{\beta}, \bar{B} \rangle \rightarrow \langle J_\beta^X, X \upharpoonright \beta, B \rangle$ be Σ_0 -elementary and $\pi[\bar{\beta}]$ be

cofinal in β . Then π is even Σ_1 -elementary.

(b) Let $\langle J_{\bar{\nu}}^X, X \upharpoonright \bar{\nu}, \bar{A} \rangle$ be rudimentary closed and $\pi : \langle J_{\bar{\nu}}^X, X \upharpoonright \bar{\nu} \rangle \rightarrow \langle J_{\bar{\nu}}^Y, Y \upharpoonright \bar{\nu} \rangle$ be Σ_0 -elementary and cofinal. Then there is a uniquely determined $A \subseteq J_{\bar{\nu}}^Y$ such that $\pi : \langle J_{\bar{\nu}}^X, X \upharpoonright \bar{\nu}, \bar{A} \rangle \rightarrow \langle J_{\bar{\nu}}^Y, X \upharpoonright \bar{\nu}, A \rangle$ is Σ_0 -elementary and $\langle J_{\bar{\nu}}^Y, X \upharpoonright \bar{\nu}, A \rangle$ is rudimentary closed.

Proof: (a) Let φ be a Σ_0 formula such that $\langle J_{\bar{\beta}}^X, X \upharpoonright \bar{\beta}, B \rangle \models (\exists z)\varphi(z, \pi(x_i))$. Since $\pi[\bar{\beta}]$ is cofinal in β , there is a $\nu \in \bar{\beta}$ such that $\langle J_{\bar{\beta}}^X, X \upharpoonright \bar{\beta}, B \rangle \models (\exists z \in S_{\pi(\nu)})\varphi(z, \pi(x_i))$. Here, the S_{ν} is defined as in lemma 10. If $\pi(S_{\nu}) = S_{\pi(\nu)}$, then $\langle J_{\bar{\beta}}^X, X \upharpoonright \bar{\beta}, B \rangle \models (\exists z \in \pi(S_{\nu}))\varphi(z, \pi(x_i))$. So, by the Σ_0 -elementarity of π , $\langle J_{\bar{\beta}}^X, X \upharpoonright \bar{\beta}, \bar{B} \rangle \models (\exists z \in S_{\nu})\varphi(z, x_i)$. I.e. $\langle J_{\bar{\beta}}^X, X \upharpoonright \bar{\beta}, \bar{B} \rangle \models (\exists z)\varphi(z, x_i)$. The converse is trivial.

It remains to prove $\pi(S_{\nu}) = S_{\pi(\nu)}$. This is done by induction on ν . If $\nu = 0$ or $\nu \notin \text{Lim}$, then the claim is obvious by the definition of S_{ν} and the induction hypothesis. So let $\lambda \in \text{Lim}$ and $M := \pi(S_{\lambda})$. Then M is transitive by the Σ_0 -elementarity of π . And since $\lambda \in \text{Lim}$ (i.e. $S_{\lambda} = J_{\lambda}^X$), $\langle S_{\nu} \mid \nu < \lambda \rangle$ is definable over $\langle J_{\lambda}^X, X \upharpoonright \lambda \rangle$ by (the proof of) lemma 10. Let φ be the formula $(\forall x)(\exists \nu)(x \in S_{\nu})$. Since π is Σ_0 -elementary, $\pi \upharpoonright S_{\lambda} : \langle J_{\lambda}^X, X \upharpoonright \lambda \rangle \rightarrow \langle M, (X \upharpoonright \lambda) \cap M \rangle$ is elementary. Thus, if $\langle J_{\lambda}^X, X \upharpoonright \lambda \rangle \models \varphi$, then also $\langle M, (X \upharpoonright \lambda) \cap M \rangle \models \varphi$. Since M is transitive, we get $M = S_{\tau}$ for a $\tau \in \text{Lim}$. And, by $\pi(\lambda) = \pi(S_{\lambda} \cap \text{On}) = S_{\tau} \cap \text{On} = \tau$, it follows that $\pi(S_{\lambda}) = S_{\pi(\lambda)}$.

(b) Since $\langle J_{\bar{\nu}}^X, X \upharpoonright \bar{\nu}, \bar{A} \rangle$ is rudimentary closed, $\bar{A} \cap S_{\mu} \in J_{\bar{\nu}}^X$ for all $\mu < \bar{\nu}$ where S_{μ} is defined as in lemma 10. As in the proof of (a), $\pi(S_{\mu}) = S_{\pi(\mu)}$. So we need $\pi(\bar{A} \cap S_{\mu}) = A \cap S_{\pi(\mu)}$ to get that $\pi : \langle J_{\bar{\nu}}^X, X \upharpoonright \bar{\nu}, \bar{A} \rangle \rightarrow \langle J_{\bar{\nu}}^Y, X \upharpoonright \bar{\nu}, A \rangle$ is Σ_0 -elementary. Since π is cofinal, we necessarily obtain $A = \bigcup \{ \pi(\bar{A} \cap S_{\mu}) \mid \mu < \bar{\nu} \}$. But then $\langle J_{\bar{\nu}}^Y, X \upharpoonright \bar{\nu}, A \rangle$ is rudimentary closed. For, if $x \in J_{\bar{\nu}}^X$, we can choose some $\mu < \bar{\nu}$ such that $x \in S_{\pi(\mu)}$. And $x \cap A = x \cap (A \cap S_{\pi(\mu)}) = x \cap \pi(\bar{A} \cap S_{\mu}) \in J_{\bar{\nu}}^X$. Now, let $\langle J_{\bar{\nu}}^X, X \upharpoonright \bar{\nu}, \bar{A} \rangle \models \varphi(x_i)$ where φ is a Σ_0 formula and $u \in J_{\bar{\nu}}^X$ is transitive such that $x_i \in u$. Then $\langle u, X \upharpoonright \bar{\nu} \cap u, A \cap u \rangle \models \varphi(x_i)$ holds. Since $\pi : \langle J_{\bar{\nu}}^X, X \upharpoonright \bar{\nu} \rangle \rightarrow \langle J_{\bar{\nu}}^Y, Y \upharpoonright \bar{\nu} \rangle$ is Σ_0 -elementary, $\langle \pi(u), Y \upharpoonright \bar{\nu} \cap \pi(u), A \cap \pi(u) \rangle \models \varphi(\pi(x_i))$. Because $\pi(u)$ is transitive, we get $\langle J_{\bar{\nu}}^Y, X \upharpoonright \bar{\nu}, A \rangle \models \varphi(\pi(x_i))$. This argument works as well for the converse. \square

Write $\text{Cond}_B(I_{\beta}^0)$ if there exists for all $H \prec_1 \langle I_{\beta}^0, B \rangle$ some $\bar{\beta}$ and some \bar{B} such that $H \cong \langle I_{\bar{\beta}}^0, \bar{B} \rangle$.

Lemma 19 (Extension of embeddings)

Let $\beta > \omega$, $m \geq 0$ and $\langle I_{\beta}^0, B \rangle$ be a rudimentary closed structure. Let $\text{Cond}_B(I_{\beta}^0)$ hold. Let ρ be the projectum of $\langle I_{\beta}^0, B \rangle$, A the standard code and p the standard parameter of $\langle I_{\beta}^0, B \rangle$. Then $\text{Cond}_A(I_{\rho}^0)$ holds. And if $\langle I_{\bar{\rho}}^0, \bar{A} \rangle$ is rudimentary closed and $\pi : \langle I_{\bar{\rho}}^0, \bar{A} \rangle \rightarrow \langle I_{\rho}^0, A \rangle$ is Σ_m -elementary, then there is a uniquely determined Σ_{m+1} -elementary extension $\tilde{\pi} : \langle I_{\bar{\beta}}^0, \bar{B} \rangle \rightarrow \langle I_{\beta}^0, B \rangle$ of π where $\bar{\rho}$ is the projectum of $\langle I_{\bar{\beta}}^0, \bar{B} \rangle$, \bar{A} is the standard code and $\tilde{\pi}^{-1}(p)$ is the standard parameter of $\langle I_{\bar{\beta}}^0, \bar{B} \rangle$.

Proof: Let $H = h_{\beta, B}[\omega \times (\text{rng}(\pi) \times \{p\})] \prec_1 \langle I_{\beta}^0, B \rangle$ and $\tilde{\pi} : \langle I_{\bar{\beta}}^0, \bar{B} \rangle \rightarrow \langle I_{\beta}^0, B \rangle$ be the uncollapse of H .

(1) $\tilde{\pi}$ is an extension of π

Let $\tilde{\rho} = \text{sup}(\pi[\bar{\rho}])$ and $\tilde{A} = A \cap J_{\tilde{\rho}}^X$. Then $\pi : \langle J_{\tilde{\rho}}^X, X \upharpoonright \tilde{\rho}, \tilde{A} \rangle \rightarrow \langle J_{\rho}^X, X \upharpoonright \rho, A \rangle$

$\tilde{\rho}, \tilde{A}$) is Σ_0 -elementary, and by lemma 18, it is even Σ_1 -elementary. We have $\text{rng}(\pi) = H \cap J_{\tilde{\rho}}^X$. Obviously $\text{rng}(\pi) \subseteq H \cap J_{\tilde{\rho}}^X$. So let $y \in H \cap J_{\tilde{\rho}}^X$. Then there is an $i \in \omega$ and an $x \in \text{rng}(\pi)$ such that y is the unique $y \in J_{\tilde{\beta}}^X$ that satisfies $\langle I_{\tilde{\beta}}^0, B \rangle \models \varphi_i(\langle y, x \rangle, p)$. So by definition of A , y is the unique $y \in J_{\tilde{\beta}}^X$ such that $\tilde{A}(i, \langle y, x \rangle)$. But $x \in \text{rng}(\pi)$ and $\pi : \langle J_{\tilde{\rho}}^X, X \upharpoonright \tilde{\rho}, \tilde{A} \rangle \rightarrow \langle J_{\tilde{\rho}}^X, X \upharpoonright \tilde{\rho}, \tilde{A} \rangle$ is Σ_1 -elementary. Therefore $y \in \text{rng}(\pi)$. So we have proved that H is an \in -end-extension of $\text{rng}(\pi)$. Since π is the collapse of $\text{rng}(\pi)$ and $\tilde{\pi}$ the collapse of H , we obtain $\pi \subseteq \tilde{\pi}$.

(2) $\tilde{\pi} : \langle I_{\tilde{\beta}}^0, \tilde{B} \rangle \rightarrow \langle I_{\tilde{\beta}}^0, B \rangle$ is Σ_{m+1} -elementary

We must prove $H \prec_{m+1} \langle I_{\tilde{\beta}}^0, B \rangle$. If $m = 0$, this is clear. So let $m > 0$ and let y be Σ_{m+1} -definable in $\langle I_{\tilde{\beta}}^0, B \rangle$ with parameters from $\text{rng}(\pi) \cup \{p\}$. Then we have to show $y \in H$. Let φ be a Σ_{m+1} formula and $x_i \in \text{rng}(\pi)$ such that y is uniquely determined by $\langle I_{\tilde{\beta}}^0, B \rangle \models \varphi(y, x_i, p)$. Let $\tilde{h}(\langle i, x \rangle) \simeq h(i, \langle x, p \rangle)$. Then $\tilde{h}[J_{\tilde{\rho}}^X] = J_{\tilde{\rho}}^X$ by the definition of p . So there is a $z \in J_{\tilde{\rho}}^X$ such that $y = \tilde{h}(z)$. If such a z lies in $J_{\tilde{\rho}}^X \cap H$, then also $y \in H$, since $z, p \in H \prec_1 \langle I_{\tilde{\beta}}^0, B \rangle$. Let $D = \text{dom}(\tilde{h}) \cap J_{\tilde{\rho}}^X$. Then it suffices to show

$$(\star) \quad (\exists z_0 \in D)(\forall z_1 \in D) \dots \langle I_{\tilde{\beta}}^0, B \rangle \models \psi(\tilde{h}(z_i), \tilde{h}(z), x_i, p)$$

for some $z \in H \cap J_{\tilde{\rho}}^X$ where ψ is Σ_1 for even m and Π_1 for odd m such that $\varphi(y, x_i, p) \Leftrightarrow \langle I_{\tilde{\beta}}^0, B \rangle \models (\exists z_0)(\forall z_1) \dots \psi(z_i, y, x_i, p)$. First, let m be even. Since A is the standard code, there is an $i_0 \in \omega$ such that $z \in D \Leftrightarrow A(i_0, x)$ holds for all $z \in J_{\tilde{\rho}}^X$ – and a $j_0 \in \omega$ such that, for all $z_i, z \in D$, $\langle I_{\tilde{\beta}}^0, B \rangle \models \psi(\tilde{h}(z_i), \tilde{h}(z), x_i, p)$ iff $A(j_0, \langle z_i, z, x_i \rangle)$. Thus (\star) is, for $z \in J_{\tilde{\rho}}^X$, equivalent with an obvious Σ_m formula. If m is odd, then write in (\star) $\dots \neg \langle I_{\tilde{\beta}}^0, B \rangle \models \neg \psi(\dots)$. Then $\neg \psi$ is Σ_1 and we can proceed as above. Eventually $\pi : \langle I_{\tilde{\rho}}^0, \tilde{A} \rangle \rightarrow \langle I_{\tilde{\rho}}^0, A \rangle$ is Σ_m -elementary by the hypothesis and $\pi \subseteq \tilde{\pi}$ by (1) – i.e. $H \cap J_{\tilde{\rho}}^X \prec_m \langle I_{\tilde{\rho}}^0, A \rangle$. Since there is a $z \in J_{\tilde{\rho}}^X$ which satisfies (\star) and $x_i, p \in H \cap J_{\tilde{\rho}}^X$, there exists such a $z \in H \cap J_{\tilde{\rho}}^X$. Let $H \prec_1 \langle I_{\tilde{\rho}}^0, A \rangle$. Let π be the uncollapse of H . Then π has a Σ_1 -elementary extension $\tilde{\pi} : \langle I_{\tilde{\beta}}^0, \tilde{B} \rangle \rightarrow \langle I_{\tilde{\beta}}^0, B \rangle$. So $H \cong \langle I_{\tilde{\rho}}^0, \tilde{A} \rangle$ for some $\tilde{\rho}$ and \tilde{A} . I.e. $\text{Cond}_A(I_{\tilde{\rho}}^0)$.

(3) $\tilde{A} = \{ \langle i, x \rangle \mid i \in \omega \text{ and } x \in J_{\tilde{\rho}}^X \text{ and } \langle I_{\tilde{\beta}}^0, \tilde{B} \rangle \models \varphi_i(x, \tilde{\pi}^{-1}(p)) \}$

Since $\pi : \langle I_{\tilde{\rho}}^0, \tilde{A} \rangle \rightarrow \langle I_{\tilde{\rho}}^0, A \rangle$ is Σ_0 -elementary, $\tilde{A}(i, x) \Leftrightarrow A(i, \pi(x))$ for $x \in J_{\tilde{\rho}}^X$. Since A is the standard code of $\langle I_{\tilde{\beta}}^0, B \rangle$, $A(i, \pi(x)) \Leftrightarrow \langle I_{\tilde{\beta}}^0, B \rangle \models \varphi_i(\pi(x), p)$. Finally, $\langle I_{\tilde{\beta}}^0, B \rangle \models \varphi_i(\pi(x), p) \Leftrightarrow \langle I_{\tilde{\beta}}^0, \tilde{B} \rangle \models \varphi_i(x, \tilde{\pi}^{-1}(p))$, because $\tilde{\pi} : \langle I_{\tilde{\beta}}^0, \tilde{B} \rangle \rightarrow \langle I_{\tilde{\beta}}^0, B \rangle$ is Σ_1 -elementary.

(4) $\tilde{\rho}$ is the projectum of $\langle I_{\tilde{\beta}}^0, \tilde{B} \rangle$

By the definition of H , $J_{\tilde{\rho}}^X = h_{\tilde{\beta}, \tilde{B}}[X \times (J_{\tilde{\rho}}^X \times \{ \tilde{\pi}^{-1}(p) \})]$. So $f(\langle i, x \rangle) \simeq h_{\tilde{\beta}, \tilde{B}}(i, \langle x, \tilde{\pi}^{-1}(p) \rangle)$ is a over $\langle I_{\tilde{\beta}}^0, \tilde{B} \rangle$ Σ_1 -definable function such that $f[J_{\tilde{\rho}}^X] = J_{\tilde{\beta}}^X$. It remains to prove that $\langle I_{\tilde{\rho}}^0, C \rangle$ is rudimentary closed for all $C \in \Sigma_1(\langle I_{\tilde{\beta}}^0, \tilde{B} \rangle) \cap \mathfrak{P}(J_{\tilde{\rho}}^X)$. By the definition of H , there exists an $i \in \omega$ and a $y \in J_{\tilde{\rho}}^X$ such that $x \in C \Leftrightarrow \langle I_{\tilde{\beta}}^0, \tilde{B} \rangle \models \varphi_i(\langle x, y \rangle, \tilde{\pi}^{-1}(p))$ for all $x \in J_{\tilde{\rho}}^X$. Thus, by (3), $x \in C \Leftrightarrow \tilde{A}(i, \langle x, y \rangle)$. For $u \in J_{\tilde{\rho}}^X$, let $v = \{ \langle i, \langle x, y \rangle \rangle \mid x \in u \}$. Then $v \in J_{\tilde{\rho}}^X$ and $\tilde{A} \cap v \in J_{\tilde{\rho}}^X$, because $\langle I_{\tilde{\rho}}^0, \tilde{A} \rangle$ is rudimentary closed by the hypothesis. But

$x \in C \cap u$ holds iff $\langle i, \langle x, y \rangle \rangle \in \bar{A} \cap v$. Finally, $J_{\bar{\rho}}^X$ is rudimentary closed and therefore $C \cap u \in J_{\bar{\rho}}^X$.

(5) $\tilde{\pi}^{-1}(p)$ is the standard parameter of $\langle I_{\bar{\beta}}^0, \bar{B} \rangle$

By the definition of H , $J_{\bar{\beta}}^X = h_{\bar{\beta}, \bar{B}}[\omega \times (J_{\bar{\rho}}^X \times \{\tilde{\pi}^{-1}(p)\})]$ and, by (4), $\bar{\rho}$ is the projectum of $\langle I_{\bar{\beta}}^0, \bar{B} \rangle$. So we just have to prove that $\tilde{\pi}^{-1}(p)$ is the least with this property. Suppose that $\bar{p}' < \tilde{\pi}^{-1}(p)$ had this property as well. Then there were an $i \in \omega$ and an $x \in J_{\bar{\rho}}^X$ such that $\tilde{\pi}^{-1}(p) = h_{\bar{\beta}, \bar{B}}(i, \langle x, \bar{p}' \rangle)$. Since $\tilde{\pi} : \langle I_{\bar{\beta}}^0, \bar{B} \rangle \rightarrow \langle I_{\beta}^0, B \rangle$ is Σ_1 -elementary, we had $p = h_{\beta, B}(i, \langle x, p' \rangle)$ for $p' = \pi(\bar{p}') < p$. And so also $h_{\beta, B}[\omega \times (J_{\rho}^X \times \{p'\})] = J_{\beta}^X$. That contradicts the definition of p .

(6) Uniqueness

Assume $\langle I_{\bar{\beta}_0}^0, \bar{B}_0 \rangle$ and $\langle I_{\bar{\beta}_1}^0, \bar{B}_1 \rangle$ both have $\bar{\rho}$ as projectum and \bar{A} as standard code. Let \bar{p}_i be the standard parameter of $\langle I_{\bar{\beta}_i}^0, \bar{B}_i \rangle$. Then, for all $j \in \omega$ and $x \in J_{\bar{\rho}}^X$, $\langle I_{\bar{\beta}_0}^0, \bar{B}_0 \rangle \models \varphi_j(x, \bar{p}_0)$ iff $\bar{A}(j, x)$ iff $\langle I_{\bar{\beta}_1}^0, \bar{B}_1 \rangle \models \varphi_j(x, \bar{p}_1)$. So $\sigma(h_{\bar{\beta}_0, \bar{B}_0}(j, \langle x, \bar{p}_0 \rangle)) \simeq h_{\bar{\beta}_1, \bar{B}_1}(j, \langle x, \bar{p}_1 \rangle)$ defines an isomorphism $\sigma : \langle I_{\bar{\beta}_0}^0, \bar{B}_0 \rangle \cong \langle I_{\bar{\beta}_1}^0, \bar{B}_1 \rangle$, because, for both i , $h_{\bar{\beta}_i, \bar{B}_i}[\omega \times (J_{\bar{\rho}}^X \times \{\bar{p}_i\})] = J_{\bar{\beta}_i}^X$ holds. But since both structures are transitive, σ must be the identity. Finally, let $\tilde{\pi}_0 : \langle I_{\bar{\beta}_0}^0, \bar{B}_0 \rangle \rightarrow \langle I_{\beta}^0, B \rangle$ and $\tilde{\pi}_1 : \langle I_{\bar{\beta}_1}^0, \bar{B}_1 \rangle \rightarrow \langle I_{\beta}^0, B \rangle$ be Σ_1 -elementary extensions of π . Let \bar{p} be the standard parameter of $\langle I_{\bar{\beta}}^0, \bar{B} \rangle$. Then, for every $y \in J_{\bar{\rho}}^X$, there is an $x \in J_{\bar{\rho}}^X$ and a $j \in \omega$ such that $y = h_{\bar{\beta}, \bar{B}}(j, \langle x, \bar{p} \rangle)$ – and $\tilde{\pi}_0(y) = h_{\beta, B}(j, \pi(x), \pi(p)) = \tilde{\pi}_1(y)$. Thus $\tilde{\pi}_0 = \tilde{\pi}_1$. \square

To code the Σ_n information of I_{β} where $\beta \in S^X$ in a structure $\langle I_{\rho}^0, A \rangle$, one iterates this process.

For $n \geq 0$, $\beta \in S^X$, let

$$\begin{aligned} \rho^0 &= \beta, p^0 = \emptyset, A^0 = X_{\beta} \\ \rho^{n+1} &= \text{the projectum of } \langle I_{\rho^n}^0, A^n \rangle \\ p^{n+1} &= \text{the standard parameter of } \langle I_{\rho^n}^0, A^n \rangle \\ A^{n+1} &= \text{the standard code of } \langle I_{\rho^n}^0, A^n \rangle. \end{aligned}$$

Call

$$\begin{aligned} \rho^n &\text{ the } n\text{-th projectum of } \beta, \\ p^n &\text{ the } n\text{-th (standard) parameter of } \beta, \\ A^n &\text{ the } n\text{-th (standard) code of } \beta. \end{aligned}$$

By lemma 17, $A^n \subseteq J_{\rho^n}^X$ is Σ_n -definable over I_{β} and, for all $m \geq 1$,

$$\Sigma_{n+m}(I_{\beta}) \cap \mathfrak{P}(J_{\rho^n}^X) = \Sigma_m(\langle I_{\rho^n}^0, A^n \rangle).$$

From lemma 19, we get by induction:

For $\beta > \omega$, $n \geq 1$, $m \geq 0$, let ρ^n be the n -th projectum and A^n be the n -th code of β . Let $\langle I_{\bar{\rho}}^0, \bar{A} \rangle$ be a rudimentary closed structure and $\pi : \langle I_{\bar{\rho}}^0, \bar{A} \rangle \rightarrow \langle I_{\rho^n}^0, A^n \rangle$ be Σ_m -elementary. Then:

(1) There is a unique $\bar{\beta} \geq \bar{\rho}$ such that $\bar{\rho}$ is the n -th projectum and \bar{A} is the n -th code of $\bar{\beta}$.

For $k \leq n$ let

ρ^k be the k -th projectum of β ,
 p^k the k -th parameter of β ,
 A^k the k -th code of β

and

$\bar{\rho}^k$ the k -th projectum of $\bar{\beta}$,
 \bar{p}^k the k -th parameter of $\bar{\beta}$,
 \bar{A}^k the k -th code of $\bar{\beta}$.

(2) There exists a unique extension $\tilde{\pi}$ of π such that, for all $0 \leq k \leq n$,

$\tilde{\pi} \upharpoonright J_{\bar{\rho}^k}^X : \langle I_{\bar{\rho}^k}^0, \bar{A}^k \rangle \rightarrow \langle I_{\rho^k}^0, A^k \rangle$ is Σ_{m+n-k} -elementary

and $\tilde{\pi}(\bar{p}^k) = p^k$.

Lemma 20

Let $\omega < \beta \in S^X$. Then all projecta of β exist.

Proof by induction on n . That ρ^0 exists is clear. So suppose that the first projecta $\rho^0, \dots, \rho^{n-1}, \rho := \rho^n$, the parameters p^0, \dots, p^n and the codes $A^0, \dots, A^{n-1}, A := A^n$ of β exist. Let $\gamma \in \text{Lim}$ be minimal such that there is some over $\langle I_\rho^0, A \rangle$ Σ_1 -definable function f such that $f[J_\gamma^X] = J_\rho^X$. Let $C \in \Sigma_1(\langle I_\rho^0, A \rangle) \cap \mathfrak{P}(J_\gamma^X)$. We have to prove that $\langle I_\gamma^0, C \rangle$ is rudimentary closed. If $\gamma = \omega$, then $J_\gamma^X = H_\omega$, and this is obvious. If $\gamma > \omega$, then $\gamma \in \text{Lim}^2$ by the definition of γ . Then it suffices to show $C \cap J_\delta^X \in J_\gamma^X$ for $\delta \in \text{Lim} \cap \gamma$. Let $B := C \cap J_\delta^X$ be definable over $\langle I_\rho^0, A \rangle$ with parameter q . Since obviously $\gamma \leq \rho$, $C \cap J_\delta^X$ is Σ_n -definable over I_β with parameters p_1, \dots, p^n, q by lemma 17. So let φ be a Σ_n formula such that $x \in C \Leftrightarrow I_\beta \models \varphi(x, p^1, \dots, p^n, q)$. Let

$$\begin{aligned} H_{n+1} &:= h_{\rho^n, A^n}[\omega \times (J_\delta^X \times \{q\})] \\ H_n &:= h_{\rho^{n-1}, A^{n-1}}[\omega \times (H_n \times \{p^n\})] \\ H_{n-1} &:= h_{\rho^{n-2}, A^{n-2}}[\omega \times (H_{n-1} \times \{p^{n-1}\})] \\ &\text{etc.} \end{aligned}$$

Since $L[X]$ has condensation, there is an I_μ such that $H_1 \cong I_\mu$. Let π be the uncollapse of H_1 . Then π is the extension of the collapse of H_{n+1} defined in the proof of lemma 19. Therefore it is Σ_{n+1} -elementary. Since $B \subseteq J_\delta^X$ and $\pi \upharpoonright J_\delta^X = id \upharpoonright J_\delta^X$, we get $x \in B \Leftrightarrow I_\mu \models \varphi(x, \pi^{-1}(p^1), \dots, \pi^{-1}(p^n), \pi^{-1}(q))$. So B is indeed already Σ_n -definable over I_μ . Thus $B \in J_{\mu+1}^X$ by lemma 8. But now we are done because $\mu < \rho$. For, if

$$\begin{aligned} h_{n+1}(\langle i, x \rangle) &= h_{\rho^n, A^n}(i, \langle x, p \rangle) \\ h_n(\langle i, x \rangle) &= h_{\rho^{n-1}, A^{n-1}}(i, \langle x, p^n \rangle) \\ &\text{etc.} \end{aligned}$$

then the function $h = h_1 \circ \dots \circ h_{n+1}$ is Σ_{n+1} -definable over I_β . Thus the function $\bar{h} = \pi[h \cap (H_1 \times H_1)]$ is Σ_{n+1} -definable over I_μ and $\bar{h}[J_\delta^X] = J_\mu^X$. So $\bar{h} \cap (J_\rho^X)^2$ is Σ_1 -definable over $\langle I_\rho^0, A \rangle$ by lemma 17 and lemma 19. And by the definition of γ , there is an over $\langle I_\rho^0, A \rangle$ Σ_1 -definable function f such that $f[J_\gamma^X] = J_\rho^X$. So if we had $\mu \geq \rho$, then $f \circ \bar{h}$ was an over $\langle I_\rho^0, A \rangle$ Σ_1 -definable function such that $(f \circ \bar{h})[J_\delta^X] = J_\rho^X$. That contradicts the minimality of γ . \square

Let $\omega < \nu \in S^X$, ρ^n the n -th projectum of ν , p^n the n -th parameter and A^n the n -th Code. Let

$$\begin{aligned} h_{n+1}(i, x) &= h_{\rho^n, A^n}(i, x) \\ h_n(\langle i, x \rangle) &= h_{\rho^{n-1}, A^{n-1}}(i, \langle x, p^n \rangle) \end{aligned}$$

etc.

Then define

$$h_\nu^{n+1} = h_1 \circ \dots \circ h_{n+1}.$$

We have:

- (1) h_ν^n is Σ_n -definable over I_ν
- (2) $h_\nu^n[\omega \times Q] \prec_n I_\nu$, if $Q \subseteq J_{\rho^{n-1}}^X$ is closed under ordered pairs.

Lemma 21

Let $\omega < \beta \in S^X$ and $n \geq 1$. Then

- (1) the least ordinal $\gamma \in Lim$ such that there is a over I_β Σ_n -definable function f such that $f[J_\gamma^X] = J_\beta^X$,
 - (2) the last ordinal $\gamma \in Lim$ such that $\langle I_\gamma^0, C \rangle$ is rudimentary closed for all $C \in \Sigma_n(I_\beta) \cap \mathfrak{P}(J_\gamma^X)$,
 - (3) the least ordinal $\gamma \in Lim$ such that $\mathfrak{P}(\gamma) \cap \Sigma_n(I_\beta) \not\subseteq J_\beta^X$,
- is the n -th projectum of β .

Proof:

(1) By the definition of the n -th projectum, there is an over $\langle I_{\rho^{n-1}}^0, A^{n-1} \rangle$ Σ_1 -definable f^n such that $f^n[J_{\rho^n}^X] = J_{\rho^{n-1}}^X$, an over $\langle I_{\rho^{n-2}}^0, A^{n-2} \rangle$ Σ_1 -definable f^{n-1} such that $f^{n-1}[J_{\rho^{n-1}}^X] = J_{\rho^{n-2}}^X$, etc. But then f^k is Σ_k -definable over I_β by lemma 17. So $f = f^1 \circ f^2 \circ \dots \circ f^n$ is Σ_n -definable over I_β and $f[J_{\rho^n}^X] = J_\beta^X$.

On the other hand, the projectum $\bar{\rho}$ of a rudimentary closed structure $\langle I_\beta^0, B \rangle$ is the least $\bar{\rho}$ such that there is an over $\langle I_{\bar{\rho}}^0, B \rangle$ Σ_1 -definable function f such that $f[J_{\bar{\rho}}^X] = J_\beta^X$. For, suppose there is no such $\rho < \bar{\rho}$ such that such an f , $f[J_\rho^X] = J_\beta^X$, exists. Then the proof of lemma 16 provides a contradiction. So if there was a $\gamma < \rho^n$ such that there is an over I_β Σ_n -definable function f such that $f[J_\gamma^X] = J_\beta^X$, then $g := f \cap (J_{\rho^{n-1}}^X)^2$ would be an over $\langle I_{\rho^{n-1}}^0, A^{n-1} \rangle$ Σ_1 -definable function such that $g[J_\gamma^X] = J_{\rho^{n-1}}^X$. But this is impossible.

(2) By the definition of the n -th projectum, $\langle I_{\rho^n}^0, C \rangle$ is rudimentary closed for all $C \in \Sigma_1(\langle I_{\rho^{n-1}}^0, A^{n-1} \rangle) \cap \mathfrak{P}(J_{\rho^n}^X)$. But by lemma 17, $\Sigma_1(\langle I_{\rho^{n-1}}^0, A^{n-1} \rangle) = \Sigma_n(I_\beta) \cap \mathfrak{P}(J_{\rho^{n-1}}^X)$. So, since $\rho^n \leq \rho^{n-1}$, $\langle I_{\rho^n}^0, C \rangle$ is rudimentary closed for all $C \in \Sigma_n(I_\beta) \cap \mathfrak{P}(J_{\rho^n}^X)$.

Assume γ were a larger ordinal $\in Lim$ having this property. Let f be, by (1), an over I_β Σ_n -definable function such that $f[J_\gamma^X] = J_\beta^X$. Set $C = \{u \in J_{\rho^n}^X \mid u \notin f(u)\}$. Then C is Σ_n -definable over I_β and $C \subseteq J_{\rho^n}^X$. So $\langle J_\gamma^X, C \rangle$ was rudimentary closed. And therefore $C = C \cap J_{\rho^n}^X \in J_\gamma^X \subseteq J_\beta^X$ and $C = f(u)$ for some $u \in J_{\rho^n}^X$. But this implies the contradiction that $u \in f(u) \Leftrightarrow u \in C \Leftrightarrow u \notin f(u)$.

(3) Let $\rho := \rho^n$ and f by (1) an over I_β Σ_n -definable function such that $f[J_\rho^X] = J_\beta^X$. Let j be an over I_ρ^0 Σ_1 -definable function from ρ onto J_ρ^X . Let $C = \{\nu \in \rho \mid \nu \notin f \circ j(\nu)\}$. Then C is an over I_β Σ_n -definable subset of ρ . If $C \in J_\beta^X$, then there would be a $\nu \in \rho$ such that $C = f \circ j(\nu)$, and we had the contradiction $\nu \in C \Leftrightarrow \nu \notin f \circ j(\nu) \Leftrightarrow \nu \notin C$. Thus $\mathfrak{P}(\rho) \cap \Sigma_n(I_\beta) \not\subseteq J_\beta^X$. But if $\gamma \in Lim \cap \rho$ and $D \in \mathfrak{P}(\gamma) \cap \Sigma_n(I_\beta)$, then $D = D \cap J_\gamma^X \in J_\rho^X \subseteq J_\beta^X$. So $\mathfrak{P}(\gamma) \cap \Sigma_n(I_\beta) \subseteq J_\beta^X$. \square

3 Morasses

Let $\omega_1 \leq \beta$, $S = \text{Lim} \cap \omega_{1+\beta}$ and $\kappa := \omega_{1+\beta}$.

We write Card for the class of cardinals and RCard for the class of regular cardinals.

Let \triangleleft be a binary relation on S such that:

- (a) If $\nu \triangleleft \tau$, then $\nu < \tau$.
 For all $\nu \in S - \text{RCard}$, $\{\tau \mid \nu \triangleleft \tau\}$ is closed.
 For $\nu \in S - \text{RCard}$, there is a largest μ such that $\nu \sqsubseteq \mu$.

Let μ_ν be this largest μ with $\nu \sqsubseteq \mu$.

Let

$$\nu \sqsubseteq \tau :\Leftrightarrow \nu \in \text{Lim}(\{\delta \mid \delta \triangleleft \tau\}) \cup \{\delta \mid \delta \sqsubseteq \tau\}.$$

- (b) \sqsubseteq is a (many-rooted) tree.

Hence, if $\nu \notin \text{RCard}$ is a successor in \sqsubseteq , then μ_ν is the largest μ such that $\nu \sqsubseteq \mu$. To see this, let μ_ν^* be the largest μ such that $\nu \sqsubseteq \mu$. It is clear that $\mu_\nu \leq \mu_\nu^*$, since $\nu \sqsubseteq \mu$ implies $\nu \sqsubseteq \mu$. So assume that $\mu_\nu < \mu_\nu^*$. Then $\nu \not\triangleleft \mu_\nu^*$ by the definition of μ_ν . Hence $\nu \in \text{Lim}(\{\delta \mid \delta \triangleleft \mu_\nu^*\})$ and $\nu \in \text{Lim}(\{\delta \mid \delta \sqsubseteq \mu_\nu^*\})$. Therefore, $\nu \in \text{Lim}(\sqsubseteq)$ since \sqsubseteq is a tree. That contradicts our assumption that ν is a successor in \sqsubseteq .

For $\alpha \in S$, let $|\alpha|$ be the rank of α in this tree. Let

$$\begin{aligned} S^+ &:= \{\nu \in S \mid \nu \text{ is a successor in } \sqsubseteq\} \\ S^0 &:= \{\alpha \in S \mid |\alpha| = 0\} \\ \widehat{S}^+ &:= \{\mu_\tau \mid \tau \in S^+ - \text{RCard}\} \\ \widehat{S} &:= \{\mu_\tau \mid \tau \in S - \text{RCard}\}. \end{aligned}$$

Let $S_\alpha := \{\nu \in S \mid \nu \text{ is a direct successor of } \alpha \text{ in } \sqsubseteq\}$. For $\nu \in S^+$, let α_ν be the direct predecessor of ν in \sqsubseteq . For $\nu \in S^0$, let $\alpha_\nu := 0$. For $\nu \notin S^+ \cup S^0$, let $\alpha_\nu := \nu$.

- (c) For $\nu, \tau \in (S^+ \cup S^0) - \text{RCard}$ such that $\alpha_\nu = \alpha_\tau$, suppose:

$$\nu < \tau \quad \Rightarrow \quad \mu_\nu < \tau.$$

For all $\alpha \in S$, suppose:

- (d) S_α is closed
- (e) $\text{card}(S_\alpha) \leq \alpha^+$
 $\text{card}(S_\alpha) \leq \text{card}(\alpha)$ if $\text{card}(\alpha) < \alpha$
- (f) $\omega_1 = \max(S^0) = \sup(S^0 \cap \omega_1)$
 $\omega_{1+i+1} = \max(S_{\omega_{1+i}}) = \sup(S_{\omega_{1+i}} \cap \omega_{1+i+1})$ for all $i < \beta$.

Let $D = \langle D_\nu \mid \nu \in \widehat{S} \rangle$ be a sequence such that $D_\nu \subseteq J_\nu^D$. To simplify matters, my definition of J_ν^D is such that $J_\nu^D \cap \text{On} = \nu$ (see section 3 or [SchZe]).

Let an $\langle S, \triangleleft, D \rangle$ -maplet f be a triple $\langle \bar{\nu}, |f|, \nu \rangle$ such that $\bar{\nu}, \nu \in S - \text{RCard}$ and $|f| : J_{\mu_{\bar{\nu}}}^D \rightarrow J_{\mu_\nu}^D$.

Let $f = \langle \bar{\nu}, |f|, \nu \rangle$ be an $\langle S, \triangleleft, D \rangle$ -maplet. Then we define $d(f)$ and $r(f)$ by $d(f) = \bar{\nu}$ and $r(f) = \nu$. Set $f(x) := |f|(x)$ for $x \in J_{\mu_{\bar{\nu}}}^D$ and $f(\mu_{\bar{\nu}}) := \mu_\nu$.

But $dom(f)$, $rng(f)$, $f \upharpoonright X$, etc. keep their usual set-theoretical meaning, i.e. $dom(f) = dom(|f|)$, $rng(f) = rng(|f|)$, $f \upharpoonright X = |f| \upharpoonright X$, etc.

For $\bar{\tau} \leq \mu_{\bar{\nu}}$, let $f^{(\bar{\tau})} = \langle \bar{\tau}, |f| \upharpoonright J_{\mu_{\bar{\tau}}}^D, \tau \rangle$ where $\tau = f(\bar{\tau})$. Of course, $f^{(\bar{\tau})}$ needs not to be a maplet. The same is true for the following definitions. Let $f^{-1} = \langle \nu, |f|^{-1}, \bar{\nu} \rangle$. For $g = \langle \nu, |g|, \nu' \rangle$ and $f = \langle \bar{\nu}, |f|, \nu \rangle$, let $g \circ f = \langle \bar{\nu}, |g| \circ |f|, \nu' \rangle$. If $g = \langle \nu', |g|, \nu \rangle$ and $f = \langle \bar{\nu}, |f|, \nu \rangle$ such that $rng(f) \subseteq rng(g)$, then set $g^{-1}f = \langle \bar{\nu}, |g|^{-1} \upharpoonright |f|, \nu' \rangle$. Finally set $id_{\nu} = \langle \nu, id \upharpoonright J_{\mu_{\nu}}^D, \nu \rangle$.

Let \mathfrak{F} be a set of $\langle S, \triangleleft, D \rangle$ -maplets $f = \langle \bar{\nu}, |f|, \nu \rangle$ such that the following holds:

- (0) $f(\bar{\nu}) = \nu$, $f(\alpha_{\bar{\nu}}) = \alpha_{\nu}$ and $|f|$ is order-preserving.
- (1) For $f \neq id_{\bar{\nu}}$, there is some $\beta \sqsubseteq \alpha_{\bar{\nu}}$ such that $f \upharpoonright \beta = id \upharpoonright \beta$ and $f(\beta) > \beta$.
- (2) If $\bar{\tau} \in S^+$ and $\bar{\nu} \sqsubset \bar{\tau} \sqsubseteq \mu_{\bar{\nu}}$, then $f^{(\bar{\tau})} \in \mathfrak{F}$.
- (3) If $f, g \in \mathfrak{F}$ and $d(g) = r(f)$, then $g \circ f \in \mathfrak{F}$.
- (4) If $f, g \in \mathfrak{F}$, $r(g) = r(f)$ and $rng(f) \subseteq rng(g)$, then $g^{-1} \circ f \in \mathfrak{F}$.

We write $f : \bar{\nu} \Rightarrow \nu$ if $f = \langle \bar{\nu}, |f|, \nu \rangle \in \mathfrak{F}$. If $f \in \mathfrak{F}$ and $r(f) = \nu$, then we write $f \Rightarrow \nu$. The uniquely determined β in (1) shall be denoted by $\beta(f)$.

Say $f \in \mathfrak{F}$ is minimal for a property $P(f)$ if $P(f)$ holds and $P(g)$ implies $g^{-1}f \in \mathfrak{F}$.

Let

$f_{(u,x,\nu)}$ = the unique minimal $f \in \mathfrak{F}$ for $f \Rightarrow \nu$ and $u \cup \{x\} \subseteq rng(f)$,

if such an f exists. The axioms of the morass will guarantee that $f_{(u,x,\nu)}$ always exists if $\nu \in S - RCard^{L_{\kappa}[D]}$. Therefore, we will always assume and explicitly mention that $\nu \in S - RCard^{L_{\kappa}[D]}$ when $f_{(u,x,\nu)}$ is mentioned.

Say $\nu \in S - RCard^{L_{\kappa}[D]}$ is independent if $d(f_{(\beta,0,\nu)}) < \alpha_{\nu}$ holds for all $\beta < \alpha_{\nu}$.

For $\tau \sqsubseteq \nu \in S - RCard^{L_{\kappa}[D]}$, say ν is ξ -dependent on τ if $f_{(\alpha_{\tau},\xi,\nu)} = id_{\nu}$.

For $f \in \mathfrak{F}$, let $\lambda(f) := sup(f[d(f)])$.

For $\nu \in S - RCard^{L_{\kappa}[D]}$ let

$$C_{\nu} = \{\lambda(f) < \nu \mid f \Rightarrow \nu\}$$

$$\Lambda(x, \nu) = \{\lambda(f_{(\beta,x,\nu)}) < \nu \mid \beta < \nu\}.$$

It will be shown that C_{ν} and $\Lambda(x, \nu)$ are closed in ν .

Recursively define a function $q_{\nu} : k_{\nu} + 1 \rightarrow On$, where $k_{\nu} \in \omega$:

$$q_{\nu}(0) = 0$$

$$q_{\nu}(k+1) = max(\Lambda(q_{\nu} \upharpoonright (k+1), \nu))$$

if $max(\Lambda(q_{\nu} \upharpoonright (k+1), \nu))$ exists. The axioms will guarantee that this recursion breaks off (see lemma 4 below), i.e. there is some k_{ν} such that either

$$\Lambda(q_{\nu} \upharpoonright (k_{\nu} + 1), \nu) = \emptyset$$

or

$$\Lambda(q_{\nu} \upharpoonright (k_{\nu} + 1), \nu) \text{ is unbounded in } \nu.$$

Define by recursion on $1 \leq n \in \omega$, simultaneously for all $\nu \in S - RCard^{L_{\kappa}[D]}$, $\beta \in \nu$ and $x \in J_{\mu_{\nu}}^D$ the following notions:

$$f_{(\beta,x,\nu)}^1 = f_{(\beta,x,\nu)}$$

$\tau(n, \nu) =$ the least $\tau \in S^0 \cup S^+ \cup \widehat{S}$ such that for some $x \in J_{\mu_\nu}^D$

$$f_{(\alpha_\tau,x,\nu)}^n = id_\nu$$

$x(n, \nu) =$ the least $x \in J_{\mu_\nu}^D$ such that $f_{(\alpha_\tau(n,\nu),x,\nu)}^n = id_\nu$

$$K_\nu^n = \{d(f_{(\beta,x(n,\nu),\nu)}^n) < \alpha_\tau(n,\nu) \mid \beta < \nu\}$$

$f \Rightarrow_n \nu$ iff $f \Rightarrow \nu$ and for all $1 \leq m < n$

$$rng(f) \cap J_{\alpha_\tau(m,\nu)}^D \prec_1 \langle J_{\alpha_\tau(m,\nu)}^D, D \upharpoonright \alpha_\tau(m,\nu), K_\nu^m \rangle$$

$$x(m, \nu) \in rng(f)$$

$f_{(u,\nu)}^n =$ the minimal $f \Rightarrow_n \nu$ such that $u \subseteq rng(f)$

$$f_{(\beta,x,\nu)}^n = f_{(\beta \cup \{x\}, \nu)}^n$$

$f : \bar{\nu} \Rightarrow_n \nu \Leftrightarrow f \Rightarrow_n \nu$ and $f : \bar{\nu} \Rightarrow \nu$.

Here definitions are to be understood in Kleene's sense, i.e., that the left side is defined iff the right side is, and in that case, both are equal.

Let

$n_\nu =$ the least n such that $f_{(\gamma,x,\mu_\nu)}^n$ is confinal in ν for some $x \in J_{\mu_\nu}^D$, $\gamma \sqsubset \nu$

$x_\nu =$ the least x such that $f_{(\alpha_\nu,x,\mu_\nu)}^{n_\nu} = id_{\mu_\nu}$.

Let

$$\alpha_\nu^* = \alpha_\nu \text{ if } \nu \in S^+$$

$$\alpha_\nu^* = \sup\{\alpha < \nu \mid \beta(f_{(\alpha,x_\nu,\mu_\nu)}^{n_\nu}) = \alpha\} \text{ if } \nu \notin S^+.$$

Let $P_\nu := \{x_\tau \mid \nu \sqsubset \tau \sqsubseteq \mu_\nu, \tau \in S^+\} \cup \{x_\nu\}$.

We say that $\mathfrak{M} = \langle S, \triangleleft, \mathfrak{F}, D \rangle$ is an (ω_1, β) -morass if the following axioms hold:

(MP – minimum principle)

If $\nu \in S - RCard^{L_\kappa[D]}$ and $x \in J_{\mu_\nu}^D$, then $f_{(0,x,\nu)}$ exists.

(LP1 – first logical preservation axiom)

If $f : \bar{\nu} \Rightarrow \nu$, then $|f| : \langle J_{\mu_{\bar{\nu}}}^D, D \upharpoonright \mu_{\bar{\nu}} \rangle \rightarrow \langle J_{\mu_\nu}^D, D \upharpoonright \mu_\nu \rangle$ is Σ_1 -elementary.

(LP2 – second logical preservation axiom)

Let $f : \bar{\nu} \Rightarrow \nu$ and $f(\bar{x}) = x$. Then

$$(f \upharpoonright J_{\bar{\nu}}^D) : \langle J_{\bar{\nu}}^D, D \upharpoonright \bar{\nu}, \Lambda(\bar{x}, \bar{\nu}) \rangle \rightarrow \langle J_\nu^D, D \upharpoonright \nu, \Lambda(x, \nu) \rangle$$

is Σ_0 -elementary.

(CP1 – first continuity principle)

For $i \leq j < \lambda$, let $f_i : \nu_i \Rightarrow \nu$ and $g_{ij} : \nu_i \Rightarrow \nu_j$ such that $g_{ij} = f_j^{-1} f_i$. Let $\langle g_i \mid i < \lambda \rangle$ be the transitive, direct limit of the directed system $\langle g_{ij} \mid i \leq j < \lambda \rangle$ and $hg_i = f_i$ for all $i < \lambda$. Then $g_i, h \in \mathfrak{F}$.

(CP2 – second continuity principle)

Let $f : \bar{\nu} \Rightarrow \nu$ and $\lambda = \text{sup}(f[\bar{\nu}])$. If, for some $\bar{\lambda}$, $h : \langle J_{\bar{\lambda}}^D, \bar{D} \rangle \rightarrow \langle J_{\lambda}^D, D \upharpoonright \lambda \rangle$ is Σ_1 -elementary and $\text{rng}(f \upharpoonright J_{\bar{\nu}}^D) \subseteq \text{rng}(h)$, then there is some $g : \bar{\lambda} \Rightarrow \lambda$ such that $g \upharpoonright J_{\bar{\lambda}}^D = h$.

(CP3 – third continuity principle)

If $C_{\nu} = \{\lambda(f) < \nu \mid f \Rightarrow \nu\}$ is unbounded in $\nu \in S - \text{RCard}^{L_{\kappa}[D]}$, then the following holds for all $x \in J_{\mu_{\nu}}^D$:

$$\text{rng}(f_{(0,x,\nu)}) = \bigcup \{\text{rng}(f_{(0,x,\lambda)}) \mid \lambda \in C_{\nu}\}.$$

(DP1 – first dependency axiom)

If $\mu_{\nu} < \mu_{\alpha_{\nu}}$, then $\nu \in S - \text{RCard}^{L_{\kappa}[D]}$ is independent.

(DP2 – second dependency axiom)

If $\nu \in S - \text{RCard}^{L_{\kappa}[D]}$ is η -dependent on $\tau \sqsubseteq \nu$, $\tau \in S^+$, $f : \bar{\nu} \Rightarrow \nu$, $f(\bar{\tau}) = \tau$ and $\eta \in \text{rng}(f)$, then $f(\bar{\tau}) : \bar{\tau} \Rightarrow \tau$.

(DP3 – third dependency axiom)

For $\nu \in \widehat{S} - \text{RCard}^{L_{\kappa}[D]}$ and $1 \leq n \in \omega$, the following holds:

- (a) If $f_{(\alpha_{\tau}, x, \nu)}^n = \text{id}_{\nu}$, $\tau \in S^+ \cup S^0$ and $\tau \sqsubseteq \nu$, then $\mu_{\nu} = \mu_{\tau}$.
- (b) If $\beta < \alpha_{\tau(n, \nu)}$, then also $d(f_{(\beta, x(n, \nu), \nu)}^n) < \alpha_{\tau(n, \nu)}$.

(DF – definability axiom)

(a) If $f_{(0, z_0, \nu)} = \text{id}_{\nu}$ for some $\nu \in \widehat{S} - \text{RCard}^{L_{\kappa}[D]}$ and $z_0 \in J_{\mu_{\nu}}^D$, then

$$\{\langle z, x, f_{(0, z, \nu)}(x) \rangle \mid z \in J_{\mu_{\nu}}^D, x \in \text{dom}(f_{(0, z, \nu)})\}$$

is uniformly definable over $\langle J_{\mu_{\nu}}^D, D \upharpoonright \mu_{\nu}, D_{\mu_{\nu}} \rangle$.

(b) For all $\nu \in S - \text{RCard}^{L_{\kappa}[D]}$, $x \in J_{\mu_{\nu}}^D$, the following holds:

$$f_{(0, x, \nu)} = f_{(0, \langle x, \nu, \alpha_{\nu}^*, P_{\nu} \rangle, \mu_{\nu})}^{n_{\nu}}.$$

This finishes the definition of an (ω_1, β) -morass.

A consequence of the axioms is (\times) by [Irr2]:

Theorem

$$\begin{aligned} & \{\langle z, \tau, x, f_{(0, z, \tau)}(x) \rangle \mid \tau < \nu, \mu_{\tau} = \nu, z \in J_{\mu_{\tau}}^D, x \in \text{dom}(f_{(0, z, \tau)})\} \\ & \cup \{\langle z, x, f_{(0, z, \nu)}(x) \rangle \mid \mu_{\nu} = \nu, z \in J_{\mu_{\nu}}^D, x \in \text{dom}(f_{(0, z, \nu)})\} \\ & \cup (\sqsubseteq \cap \nu^2) \end{aligned}$$

is for all $\nu \in S$ uniformly definable over $\langle J_{\nu}^D, D \upharpoonright \nu, D_{\nu} \rangle$.

A structure $\mathfrak{M} = \langle S, \triangleleft, \mathfrak{F}, D \rangle$ is called an $\omega_{1+\beta}$ -standard morass if it satisfies all axioms of an (ω_1, β) -morass except **(DF)** which is replaced by:

$\nu \triangleleft \tau \Rightarrow \nu$ is regular in J_τ^D

and there are functions $\sigma_{(x,\nu)}$ for $\nu \in \widehat{S}$ and $x \in J_\nu^D$ such that:

(MP)⁺

$$\sigma_{(x,\nu)}[\omega] = \text{rng}(f_{(0,x,\nu)})$$

(CP1)⁺

If $f : \bar{\nu} \Rightarrow \nu$ and $f(\bar{x}) = x$, then $\sigma_{(x,\nu)} = f \circ \sigma_{(\bar{x},\bar{\nu})}$.

(CP3)⁺

If C_ν is unbounded in ν , then $\sigma_{(x,\nu)} = \bigcup \{ \sigma_{(x,\lambda)} \mid \lambda \in C_\nu, x \in J_\lambda^D \}$.

(DF)⁺

(a) If $f_{(0,x,\nu)} = \text{id}_\nu$ for some $x \in J_\nu^D$, then

$$\{ \langle i, z, \sigma_{(z,\nu)}(i) \rangle \mid z \in J_\nu^D, i \in \text{dom}(\sigma_{(z,\nu)}) \}$$

is uniformly definable over $\langle J_{\mu_\nu}^D, D \upharpoonright \mu_\nu, D_{\mu_\nu} \rangle$.

(b) If C_ν is unbounded in ν , then $D_\nu = C_\nu$. If it is bounded, then $D_\nu = \{ \langle i, \sigma_{(q_\nu,\nu)}(i) \rangle \mid i \in \text{dom}(\sigma_{(q_\nu,\nu)}) \}$.

Now, I am going to construct a κ -standard morass.

Let $\beta(\nu)$ be the least β such that $J_{\beta+1}^X \models \nu$ singular.

Let $L_\kappa[X]$ satisfy amenability, condensation and coherence such that $S^X = \{ \beta(\nu) \mid \nu \text{ singular in } L_\kappa[X] \}$ and $\text{Card}^{L_\kappa[X]} = \text{Card} \cap \kappa$.

Let

$$\nu \triangleleft \tau \quad :\Leftrightarrow \quad \nu \text{ regular in } I_\tau.$$

Let

$$E = \text{Lim} - \text{RCard}^{L_\kappa[X]}.$$

For $\nu \in E$, let

$\beta(\nu)$ = the least β such that there is a cofinal $f : a \rightarrow \nu \in \text{Def}(I_\beta)$ and $a \subseteq \nu' < \nu$

$n(\nu)$ = the least $n \geq 1$ such that such an f is Σ_n -definable over $I_{\beta(\nu)}$

$\rho(\nu)$ = the $(n(\nu) - 1)$ -th projectum of $I_{\beta(\nu)}$

A_ν = the $(n(\nu) - 1)$ -th standard code of $I_{\beta(\nu)}$

$\gamma(\nu)$ = the $n(\nu)$ -th projectum of $I_{\beta(\nu)}$.

If $\nu \in S^+ - \text{Card}$, then the $n(\nu)$ -th projectum γ of $\beta(\nu)$ is less or equal $\alpha_\nu :=$ the largest cardinal in I_ν : Since α_ν is the largest cardinal in I_ν , there is, by definition of $\beta(\nu)$ and $n(\nu)$, some over $I_{\beta(\nu)}$ $\Sigma_{n(\nu)}$ -definable function f such that $f[\alpha_\nu]$ is cofinal in ν . But, since ν is regular in $\beta(\nu)$, f cannot be an element of $J_{\beta(\nu)}^X$. So $\mathfrak{P}(\nu \times \nu) \cap \Sigma_{n(\nu)}(I_{\beta(\nu)}) \not\subseteq J_{\beta(\nu)}^X$. By lemma 14, also $\mathfrak{P}(\nu) \cap \Sigma_{n(\nu)}(I_{\beta(\nu)}) \not\subseteq J_{\beta(\nu)}^X$. Using lemma 21 (3), we get $\gamma \leq \nu$. I.e. there is an over $I_{\beta(\nu)}$ $\Sigma_{n(\nu)}$ -definable function g such that $g[\nu] = J_{\beta(\nu)}^X$. On the other hand, there is, for every $\tau < \nu$ in J_ν^X , a surjection from α_ν onto τ , because α_ν is the largest cardinal in I_ν . Let f_τ be the $<_\nu$ -least such. Define $j_1(\sigma, \tau) = f_{f(\tau)}(\sigma)$ for $\sigma, \tau < \nu$. Then j_1 is $\Sigma_{n(\nu)}$ -definable over $I_{\beta(\nu)}$ and $j_1[\alpha_\nu \times \alpha_\nu] = \nu$. By lemma 15, we obtain an

over $I_{\beta(\nu)} \Sigma_{n(\nu)}$ -definable function j_2 from a subset of α_ν onto ν . Thus $g \circ j_2$ is an over $I_{\beta(\nu)} \Sigma_{n(\nu)}$ -definable map such that $g \circ j_2[\alpha_\nu] = J_{\beta(\nu)}^X$.

Moreover, $\alpha_\nu < \nu \leq \rho(\nu)$: By definition of $\rho(\nu)$, there is an over $I_{\beta(\nu)} \Sigma_{n(\nu)-1}$ -definable function f such that $f[\rho(\nu)] = \beta(\nu)$ if $n(\nu) > 1$. But ν is $\Sigma_{n(\nu)-1}$ -regular over $I_{\beta(\nu)}$. Thus $\nu \leq \rho(\nu)$. If $n(\nu) = 1$, then $\rho(\nu) = \beta(\nu) \geq \nu$.

By the first inequality, there is a q such that every $x \in J_{\rho(\nu)}^X$ is Σ_1 -definable in $\langle I_{\rho(\nu)}^0, A_\nu \rangle$ with parameters from $\alpha_\nu \cup \{q\}$. Let p_ν be the $<_{\rho(\nu)}$ -least such.

Obviously, $p_\tau \leq p_\nu$ if $\nu \sqsubseteq \tau \sqsubseteq \mu_\nu$.

Thus $P_\nu := \{p_\tau \mid \nu \sqsubseteq \tau \sqsubseteq \mu_\nu, \tau \in S^+\}$ is finite.

Now, let $\nu \in E - S^+$. By definition of $\beta(\nu)$, there exists no cofinal $f : a \rightarrow \nu$ in J_β^X such that $a \subseteq \nu' < \nu$. So $\mathfrak{P}(\nu \times \nu) \cap \Sigma_{n(\nu)}(I_{\beta(\nu)}) \not\subseteq J_{\beta(\nu)}^X$. Then, by lemma 14, $\mathfrak{P}(\nu) \cap \Sigma_{n(\nu)}(I_{\beta(\nu)}) \not\subseteq J_{\beta(\nu)}^X$. Hence, by lemma 21 (3),

$$\gamma(\nu) \leq \nu.$$

Assume $\rho(\nu) < \nu$. Then there was an over $I_{\beta(\nu)} \Sigma_{n(\nu)-1}$ -definable f such that $f[\rho(\nu)] = \nu$. But this contradicts the definition of $n(\nu)$. So

$$\nu \leq \rho(\nu).$$

Using lemma 21 (1), it follows from the first inequality that there is some over $I_{\beta(\nu)} \Sigma_{n(\nu)}$ -definable function f such that $f[J_\nu^X] = J_{\beta(\nu)}^X$. So there is a $p \in J_{\rho(\nu)}^X$ such that every $x \in J_{\rho(\nu)}^X$ is Σ_1 -definable in $\langle I_{\rho(\nu)}^0, A_\nu \rangle$ with parameters from $\nu \cup \{p\}$. Let p_ν be the least such.

Let

$$\alpha_\nu^* = \sup\{\alpha < \nu \mid h_{\rho(\nu), A_\nu}[\omega \times (J_\alpha^X \times \{p_\nu\})] \cap \nu = \alpha\}.$$

Then $\alpha_\nu^* < \nu$ because, by definition of $\beta(\nu)$, there exists a $\nu' < \nu$ and a $p \in J_{\rho(\nu)}^X$ such that $h_{\rho(\nu), A_\nu}[\omega \times (J_{\nu'}^X \times \{p\})] \cap \nu$ is cofinal in ν . But p is in $h_{\rho(\nu), A_\nu}[\omega \times (J_\nu^X \times \{p_\nu\})]$. So there is an $\alpha < \nu$ such that $h_{\rho(\nu), A_\nu}[\omega \times (J_\alpha^X \times \{p_\nu\})] \cap \nu$ is cofinal in ν . Thus $\alpha_\nu^* < \alpha < \nu$.

If $\nu \in S^+$, then we set $\alpha_\nu^* := \alpha_\nu$.

For $\nu \in E$, let $f : \bar{\nu} \Rightarrow \nu$ iff, for some f^* ,

- (1) $f = \langle \bar{\nu}, f^* \upharpoonright J_{\mu_{\bar{\nu}}}^D, \nu \rangle$,
- (2) $f^* : I_{\mu_{\bar{\nu}}} \rightarrow I_{\mu_\nu}$ is $\Sigma_{n(\nu)}$ -elementary,
- (3) $\alpha_\nu^*, p_\nu, \alpha_{\mu_\nu}^*, P_\nu \in \text{rng}(f^*)$,
- (4) $\nu \in \text{rng}(f^*)$ if $\nu < \mu_\nu$,
- (5) $f(\bar{\nu}) = \nu$ and $\bar{\nu} \in S^+ \Leftrightarrow \nu \in S^+$.

By this, \mathfrak{F} is defined.

Set $D = X$.

Let P_ν^* be minimal such that $h_{\mu_\nu}^{n(\nu)-1}(i, P_\nu^*) = P_\nu$ for an $i \in \omega$.

Let $\alpha_{\mu_\nu}^{**}$ be minimal such that $h_{\mu_\nu}^{n(\nu)-1}(i, \alpha_{\mu_\nu}^{**}) = \alpha_{\mu_\nu}^*$ for some $i \in \omega$.

Set

$$\nu^* = \emptyset \text{ if } \nu = \rho(\nu)$$

$$\nu^* = \nu \text{ if } \nu < \rho(\nu).$$

For $\tau \in On$, let S_τ be defined as in lemma 10. For $\tau \in On$, $E_i \subseteq S_\tau$ and a Σ_0 formula φ , let

$h_{\tau, E_i}^\varphi(x_1, \dots, x_m)$ the least $x_0 \in S_\tau$ w.r.t. the canonical well-ordering such that $\langle S_\tau, E_i \rangle \models \varphi(x_i)$ if such an element exists,

and

$$h_{\tau, E_i}^\varphi(x_1, \dots, x_m) = \emptyset \text{ else.}$$

For $\tau \in On$ such that $\nu^*, \alpha_\nu^*, p_\nu, \alpha_{\mu_\nu}^{**}, P_\nu^* \in S_\tau$, let $H_\nu(\alpha, \tau)$ be the closure of $S_\alpha \cup \{\nu^*, \alpha_\nu^*, p_\nu, \alpha_{\mu_\nu}^{**}, P_\nu^*\}$ under all $h_{\tau, X \cap S_\tau, A_\nu \cap S_\tau}^\varphi$. Then $H_\nu(\alpha, \tau) \prec_1 \langle S_\tau, X \cap S_\tau, A_\nu \cap S_\tau, \{\nu^*, \alpha_\nu^*, p_\nu, \alpha_{\mu_\nu}^{**}, P_\nu^*\} \rangle$ by the definition of $h_{\tau, X \cap S_\tau, A_\nu \cap S_\tau}^\varphi$. Let $M_\nu(\alpha, \tau)$ be the collapse of $H_\nu(\alpha, \tau)$. Let τ_0 be the minimal τ such that $\nu^*, \alpha_\nu^*, p_\nu, \alpha_{\mu_\nu}^{**}, P_\nu^* \in S_\tau$. Define by induction for $\tau_0 \leq \tau < \rho(\nu)$:

$$\begin{aligned} \alpha(\tau_0) &= \alpha_\nu \\ \alpha(\tau + 1) &= \text{sup}(M_\nu(\alpha(\tau), \tau + 1) \cap \nu) \\ \alpha(\lambda) &= \text{sup}\{\alpha(\tau) \mid \tau < \lambda\} \text{ if } \lambda \in \text{Lim.} \end{aligned}$$

Set

$$\begin{aligned} B_\nu &= \{\langle \alpha(\tau), M_\nu(\alpha(\tau), \tau) \rangle \mid \tau_0 < \tau \in \rho(\nu)\} \text{ if } \nu < \rho(\nu), \\ B_\nu &= \{0\} \times A_\nu \cup \{1, \nu^*, \alpha_\nu^*, p_\nu, \alpha_{\mu_\nu}^{**}, P_\nu^*\} \text{ else.} \end{aligned}$$

Lemma 22

$B_\nu \subseteq J_\nu^X$ and $\langle I_\nu^0, B_\nu \rangle$ is rudimentary closed.

Proof: If $\nu = \rho(\nu)$, then both claims are clear. Otherwise, we first prove $M_\nu(\alpha, \tau) \in J_\nu^X$ for all $\alpha < \nu$ and all $\tau \in \rho(\nu)$ such that $\tau_0 \leq \tau < \rho(\nu)$. Let such a τ be given and $\tau' \in \rho(\nu) - \text{Lim}$ be such that $X \cap S_\tau, A_\nu \cap S_\tau \in S_{\tau'}$ (rudimentary closedness of $\langle I_{\rho(\nu)}^0, A_\nu \rangle$). Let $\eta := \text{sup}(\tau' \cap \text{Lim})$. Let H be the closure of $\alpha \cup \{\nu^*, \alpha_\nu^*, p_\nu, \alpha_{\mu_\nu}^{**}, P_\nu^*, X \cap S_\tau, S_\tau, A_\nu \cap S_\tau, \eta\}$ under all $h_{\tau'}^\varphi$. Let $\sigma : H \cong S$ be the collapse of H and $\sigma(\eta) = \bar{\eta}$. If $\eta \in S^X$, then $S = S_{\tau'}$ for some $\bar{\tau}'$ by the condensation property of $L[X]$. If $\eta \notin S^X$, then $S = S_{\bar{\tau}' \upharpoonright \bar{\eta}}$ for some $\bar{\tau}'$ where $S_{\bar{\tau}' \upharpoonright \bar{\eta}}$ is defined like $S_{\tau'}$ with $X \upharpoonright \bar{\eta}$ instead of X . The reason is that, even if $\eta \notin S^X$, it is the supremum of points in S^X , because $S^X = \{\beta(\nu) \mid \nu \text{ singular in } L_\kappa[X]\}$. In both cases, $S \in J_\nu^X$ and there is a function in $I_{\bar{\eta} + \omega}$ that maps $\alpha \cup \{\sigma(\nu^*), \sigma(\alpha_\nu^*), \sigma(p_\nu), \sigma(\alpha_{\mu_\nu}^{**}), \sigma(P_\nu^*), \sigma(X \cap S_\tau), \sigma(S_\tau), \sigma(A_\nu \cap S_\tau), \sigma(\eta)\}$ onto S . So ν would be singular in $J_{\rho(\nu)}^X$ if $\nu \leq \bar{\tau}'$. But this contradicts the definition of $\beta(\nu)$. Therefore, $\sigma(\nu^*), \sigma(\alpha_\nu^*), \sigma(p_\nu), \sigma(\alpha_{\mu_\nu}^{**}), \sigma(P_\nu^*), \sigma(X \cap S_\tau), \sigma(S_\tau), \sigma(A_\nu \cap S_\tau), \sigma(\eta) \in J_\nu^X$. Let $\bar{H}_\nu(\alpha, \tau)$ be the closure of $S_\alpha \cup \{\sigma(\nu^*), \sigma(\alpha_\nu^*), \sigma(p_\nu), \sigma(\alpha_{\mu_\nu}^{**}), \sigma(P_\nu^*), \sigma(X \cap S_\tau), \sigma(S_\tau), \sigma(A_\nu \cap S_\tau), \sigma(\eta)\}$ under all $h_{\sigma(S_\tau), \sigma(X \cap S_\tau), \sigma(A_\nu \cap S_\tau)}^\varphi$ where these are defined like h_{τ, E_i}^φ but with $\sigma(S_\tau)$ instead of S_τ . Then $\bar{H}_\nu(\alpha, \tau) \prec_1 \langle \sigma(S_\tau), \sigma(X \cap S_\tau), \sigma(A_\nu \cap S_\tau), \{\sigma(\nu^*), \sigma(\alpha_\nu^*), \sigma(p_\nu), \sigma(\alpha_{\mu_\nu}^{**}), \sigma(P_\nu^*), \sigma(X \cap S_\tau), \sigma(S_\tau), \sigma(A_\nu \cap S_\tau), \sigma(\eta)\} \rangle$ and $M_\nu(\alpha, \tau)$ is the collapse of $\bar{H}_\nu(\alpha, \tau)$. Since $\nu < \rho(\nu)$ and ν is a cardinal in $I_{\beta(\nu)}$, $J_\nu^X \models ZF^-$. So we can form the collapse inside J_ν^X . Thus $M_\nu(\alpha, \tau) \in J_\nu^X$.

Now, we turn to rudimentary closedness. Since B_ν is unbounded in ν , it suffices to prove that the initial segments of B_ν are elements of J_ν^X . Such an initial segment is of the form $\langle M_\nu(\alpha(\tau), \tau) \mid \tau < \gamma \rangle$ where $\gamma < \rho(\nu)$, and we have $H_\nu(\alpha(\tau), \delta_\tau) = H_\nu(\alpha(\tau), \tau)$ where δ_τ is for $\tau < \gamma$ the least $\eta \geq \tau$ such that $\eta \in H_\nu(\alpha(\tau), \gamma) \cup \{\gamma\}$. Since $\delta_\tau \in H_\nu(\alpha(\tau), \gamma) \prec_1 \langle S_\gamma, X \cap S_\gamma, A_\nu \cap S_\gamma, \{\dots\} \rangle$,

$(H_\nu(\alpha(\tau), \delta_\tau))^{H_\nu(\alpha(\gamma), \gamma)} = H_\nu(\alpha(\tau), \tau)$. Let $\pi : M_\nu(\alpha(\gamma), \gamma) \rightarrow S_\gamma$ be the uncollapse of $H_\nu(\alpha(\gamma), \gamma)$. Then, by the Σ_1 -elementarity of π , $M_\nu(\alpha(\tau), \tau) = M_\nu(\alpha(\tau), \delta_\tau)$ is the collapse of $(H(\alpha(\tau), \pi^{-1}(\delta_\tau)))^{M_\nu(\alpha(\gamma), \gamma)}$. So $\langle M_\nu(\alpha(\tau), \tau) \mid \tau < \gamma \rangle$ is definable from $M_\nu(\alpha(\gamma), \gamma) \in J_\nu^X$. \square

Lemma 23

For $x, y_i \in J_\nu^X$, the following are equivalent:

- (i) x is Σ_1 -definable in $\langle I_{\rho(\nu)}^0, A_\nu \rangle$ with the parameters $y_i, \nu^*, \alpha_\nu^*, p_\nu, \alpha_{\mu_\nu}^{**}, P_\nu^*$.
- (ii) x is Σ_1 -definable in $\langle I_\nu^0, B_\nu \rangle$ with the parameters y_i .

Proof: For $\nu = \rho(\nu)$, this is clear. Otherwise, let first x be uniquely determined in $\langle I_{\rho(\nu)}^0, A_\nu \rangle$ by $(\exists z)\psi(z, x, \langle y_i, \nu^*, \alpha_\nu^*, p_\nu, \alpha_{\mu_\nu}^{**}, P_\nu^* \rangle)$ where ψ is a Σ_0 formula. That is equivalent to $(\exists \tau)(\exists z \in S_\tau)\psi(z, x, \langle y_i, \nu^*, \alpha_\nu^*, p_\nu, \alpha_{\mu_\nu}^{**}, P_\nu^* \rangle)$ and that again to $(\exists \tau)H_\nu(\alpha(\tau), \tau) \models (\exists z)\psi(z, x, \langle y_i, \nu^*, \alpha_\nu^*, p_\nu, \alpha_{\mu_\nu}^{**}, P_\nu^* \rangle)$. If τ is large enough, the y_i are not moved by the collapsing map, since then $y_i \in J_{\alpha(\tau)}^X \subseteq H_\nu(\alpha(\tau), \tau)$. Let $\bar{\nu}, \alpha, p, \alpha', P$ be the images of $\nu^*, \alpha_\nu^*, p_\nu, \alpha_{\mu_\nu}^{**}, P_\nu^*$ under the collapse. Then $(\exists \tau)(y_i \in J_{\alpha(\tau)}^X \text{ and } M_\nu(\alpha(\tau), \tau) \models (\exists z)\psi(z, x, \langle y_i, \bar{\nu}, \alpha, p, \alpha', P \rangle))$ defines x . So it is definable in $\langle I_\nu^0, B_\nu \rangle$.

Since B_ν and the satisfaction relation of $\langle I_\nu^0, B \rangle$ are Σ_1 -definable over $\langle I_{\rho(\nu)}^0, A_\nu \rangle$, the converse is clear. \square

Lemma 24

Let $H \prec_1 \langle I_\nu^0, B_\nu \rangle$ for a $\nu \in E$ and $\pi : \langle I_\mu^0, B \rangle \rightarrow \langle I_\nu^0, B_\nu \rangle$ be the uncollapse of H . Then $\mu \in E$ and $B = B_\mu$.

Proof: First, we extend π like in lemma 19. Let

$$M = \{x \in J_{\rho(\nu)}^X \mid x \text{ is } \Sigma_1\text{-definable in } \langle I_{\rho(\nu)}^0, A_\nu \rangle \text{ with parameters from } \text{rng}(\pi) \cup \{p_\nu, \nu^*, \alpha_\nu^*, \alpha_{\mu_\nu}^{**}, P_\nu^*\}\}.$$

Then $\text{rng}(\pi) = M \cap J_\nu^X$. For, if $x \in M \cap J_\nu^X$, then there are by definition of M $y_i \in \text{rng}(\pi)$ such that x is Σ_1 -definable in $\langle I_{\rho(\nu)}^0, A_\nu \rangle$ with the parameters y_i and $p_\nu, \nu^*, \alpha_\nu^*, \alpha_{\mu_\nu}^{**}, P_\nu^*$. Thus it is Σ_1 -definable in $\langle I_\nu^0, B_\nu \rangle$ with the y_i by lemma 23. Therefore, $x \in \text{rng}(\pi)$ because $y_i \in \text{rng}(\pi) \prec_1 \langle I_\nu^0, B_\nu \rangle$. Let $\hat{\pi} : \langle I_\rho^0, A \rangle \rightarrow \langle I_{\rho(\nu)}^0, A_\nu \rangle$ be the uncollapse of M . Then $\hat{\pi}$ is an extension of π , since $M \cap J_\nu^X$ is an \in -initial segment of M and $\text{rng}(\pi) = M \cap J_\nu^X$. In addition, there is by lemma 19 a $\Sigma_{n(\nu)}$ -elementary extension $\tilde{\pi} : I_\beta \rightarrow I_{\beta(\nu)}$ such that ρ is the $(n(\nu) - 1)$ -th projectum of I_β and A is the $(n(\nu) - 1)$ -th standard code of it. Let $\tilde{\pi}(p) = p_\nu$ and $\tilde{\pi}(\alpha) = \alpha_\nu^*$. And we have $\tilde{\pi}(\mu) = \nu$ if $\nu < \beta(\nu)$. In this case, $\nu \in \text{rng}(\pi)$ by the definition of ν^* . Since $\tilde{\pi}$ is Σ_1 -elementary, cardinals of J_μ^X are mapped on cardinals of J_ν^X .

Assume $\nu \in S^+$. Suppose there was a cardinal $\tau > \alpha$ of J_μ^X . Then $\pi(\tau) > \alpha_\tau$ was a cardinal in J_ν^X . But this is a contradiction.

Next, we note that μ is $\Sigma_{n(\nu)}$ -singular over I_β . If $\nu \in S^+$, then, by the definition of p_ν , $J_\rho^X = h_{\rho, A}[\omega \times (\alpha \times \{p\})]$ is clear. So there is an over $\langle I_\rho^0, A \rangle$ Σ_1 -definable function from α cofinal into μ . But since ρ is the $(n(\nu) - 1)$ -th projectum and A is the $(n(\nu) - 1)$ -th code of it, this function is Σ_n -definable over I_β . Now, suppose $\nu \notin S^+$. Let $\lambda := \text{sup}(\pi[\mu])$. Since $\lambda > \alpha_\nu^*$, there is a $\gamma < \lambda$ such that

$$\text{sup}(h_{\rho(\nu), A_\nu}[\omega \times (J_\gamma^X \times \{q_\nu\})] \cap \nu) \geq \lambda.$$

And since $rng(\pi)$ is cofinal in λ , there is such a $\gamma \in rng(\pi)$. Let $\gamma = \pi(\bar{\gamma})$. By the Σ_1 -elementarity of $\tilde{\pi}$, $\bar{\gamma} < \mu$ and setting $\tilde{\pi}(q) = q_\nu$ we have for every $\eta < \mu$

$$\langle I_\rho, A \rangle \models (\exists x \in J_\gamma^X)(\exists i)h_{\rho,A}(i, \langle x, p \rangle) > \eta.$$

Hence $h_{\rho,A}[\omega \times (J_\gamma^X \times \{q\})]$ is cofinal in μ . This shows $\mu \in E$.

On the other hand, μ is $\Sigma_{n(\nu)-1}$ -regular over I_β if $n(\nu) > 1$. Assume there was an over I_β $\Sigma_{n(\nu)-1}$ -definable function f and some $x \in \mu$ such that $f[x]$ was cofinal in μ . I.e. $(\forall y \in \mu)(\exists z \in x)(f(z) > y)$ would hold in I_β . Over I_β , $(\exists z \in x)(f(z) > y)$ is $\Sigma_{n(\nu)-1}$. So it is Σ_0 over $\langle I_\rho^0, A \rangle$. But then also $(\forall y \in \mu)(\exists z \in x)(f(z) > y)$ is Σ_0 over $\langle I_\rho^0, A \rangle$ if $\mu < \rho$. Hence it is $\Sigma_{n(\nu)}$ over I_β . But then the same would hold for $\tilde{\pi}(x)$ in $I_{\beta(\nu)}$. This contradicts the definition of $n(\nu)$! Now, let $\mu = \rho$. Since α is the largest cardinal in I_μ , we had in f also an over I_β $\Sigma_{n(\nu)-1}$ -definable function from α onto ρ and therefore one from α onto β . But this contradicts lemma 21 and the fact that ρ is the $(n(\nu) - 1)$ -th projectum of β . If $n(\nu) = 1$, then we get with the same argument that μ is regular in I_β .

The previous two paragraphs show $\beta = \beta(\mu)$ and $n(\mu) = n(\nu)$. We are done if we can also show that $\alpha = \alpha_\mu^*$, $\pi(\alpha_{\mu\mu}^{**}) = \alpha_{\mu\nu}^{**}$, $p = p_\mu$, $\pi(P_\mu^*) = P_\nu^*$, because $\tilde{\pi}$ is Σ_1 -elementary, $\tilde{\pi}(h_{\tau, X \cap S_\tau, A_\mu \cap S_\tau}^\varphi(x_i)) = h_{\tilde{\pi}(\tau), X \cap S_{\tilde{\pi}(\tau)}, A_\nu \cap S_{\tilde{\pi}(\tau)}}^\varphi(x_i)$ for all Σ_1 formulas φ and $x_i \in S_\tau$.

For $\nu \in S^+$, $\alpha = \alpha_\mu$ was shown above. So let $\nu \notin S^+$. By the Σ_1 -elementarity of $\tilde{\pi}$, we have for all $\alpha \in \mu$

$$h_{\rho,A}[\omega \times (J_\alpha^X \times \{p\})] \cap \mu = \alpha \Leftrightarrow h_{\rho(\nu), A_\nu}[\omega \times (J_{\pi(\alpha)}^X \times \{p_\nu\})] \cap \nu = \pi(\alpha).$$

The same argument proves $\pi(\alpha_{\mu\mu}^{**}) = \alpha_{\mu\nu}^{**}$. Finally, $p = p_\mu$ and $\pi(P_\mu^*) = P_\nu^*$ can be shown as in (5) in the proof of lemma 19. \square

Lemma 25

Let $H \prec_1 \langle I_\nu^0, B_\nu \rangle$ and $\lambda = \sup(H \cap \nu)$ for a $\nu \in E$. Then $\lambda \in E$ and $B_\nu \cap J_\lambda^X = B_\lambda$.

Proof: Let $\pi_0 : \langle I_\mu^0, B_\mu \rangle \rightarrow \langle I_\lambda^0, B_\nu \cap J_\lambda^X \rangle$ be the uncollapse of H and let $\pi_1 : \langle I_\lambda^0, B_\nu \cap J_\lambda^X \rangle \rightarrow \langle I_\nu^0, B_\nu \rangle$ be the identity. Since $L[X]$ has coherence, π_0 and π_1 are Σ_0 -elementary. By lemma 18, π_0 is even Σ_1 -elementary, because it is cofinal. To show $B_\lambda = B_\nu \cap J_\lambda^X$, we extend π_0 and π_1 to $\hat{\pi}_0 : \langle I_{\rho(\mu)}^0, A_\mu \rangle \rightarrow \langle I_\rho^0, A \rangle$ and $\hat{\pi}_1 : \langle I_\rho^0, A \rangle \rightarrow \langle I_{\rho(\nu)}^0, A_\nu \rangle$ in such a way that $\hat{\pi}_0$ is Σ_1 -elementary and $\hat{\pi}_1$ is Σ_0 -elementary. Then we know from lemma 19 that ρ is the $(n(\nu) - 1)$ -th projectum of some β and A is the $(n(\nu) - 1)$ -th code of it. So there is a $\Sigma_{n(\nu)}$ -elementary extension of $\hat{\pi}_0 : I_{\bar{\beta}} \rightarrow I_\beta$. We can again use the argument from lemma 24 to show that λ is $\Sigma_{n(\nu)-1}$ -regular over I_β . But on the other hand, λ is as supremum of $H \cap On$ $\Sigma_{n(\nu)}$ -singular over I_β . From this, we conclude as in the proof of lemma 24 that $B_\lambda = B_\nu \cap J_\lambda^X$.

First, suppose $\nu \in S^+$. Since $\alpha_\nu \in H \prec_1 \langle I_\nu^0, B_\nu \rangle$, $\alpha_\nu < \lambda \leq \nu$. Since $I_\nu \models (\alpha_\nu \text{ is the largest cardinal})$, we therefore have $\lambda \notin Card$. In addition, α_ν is the largest cardinal in I_λ . Assume τ was the next larger cardinal. Then τ was Σ_1 -definable in I_λ with parameter α_ν and some $\tau' \in H$ and hence it was in H . By the Σ_1 -elementarity of π_0 , $\pi_0^{-1}(\tau) > \pi_0^{-1}(\alpha_\nu) = \alpha_\mu$ was also a cardinal in I_μ . But this contradicts the definition of α_μ .

But now to $B_\lambda = B_\nu \cap J_\lambda^X$. First, assume $\nu \notin S^+$. Let $\pi = \pi_1 \circ \pi_0 : \langle I_\mu^0, B_\mu \rangle \rightarrow \langle I_\nu^0, B_\nu \rangle$ and $\hat{\pi} : \langle I_{\rho(\mu)}^0, A_\mu \rangle \rightarrow \langle I_{\rho(\nu)}^0, A_\nu \rangle$ be the extension constructed in the proof of lemma 24. Let $\gamma = \text{sup}(\text{rng}(\hat{\pi}))$. Then $\hat{\pi}' = \hat{\pi} \cap (J_{\rho(\mu)}^X \times J_\gamma^X) : \langle I_{\rho(\mu)}^0, A_\mu \rangle \rightarrow \langle I_\gamma^0, A_\nu \cap J_\gamma^X \rangle$ is Σ_0 -elementary, by coherence of $L_\kappa[X]$, and cofinal. Thus $\hat{\pi}'$ is Σ_1 -elementary. Let $H' = h_{\gamma, A_\nu \cap J_\gamma^X}[\omega \times (J_\lambda^X \times \{p_\nu\})]$ and $\hat{\pi}_1 : \langle I_\rho^0, A \rangle \rightarrow \langle I_{\rho(\nu)}^0, A_\nu \rangle$ be the uncollapse of H' . Then $H = \text{rng}(\hat{\pi}') \subseteq H'$. To see this, let $z \in \text{rng}(\hat{\pi}')$ and $z = \hat{\pi}'(y)$. Then by definition of p_μ , there is an $x \in J_\mu^X$ and an $i \in \omega$ such that $y = h_{\rho(\mu), A_\mu}(i, \langle x, p_\mu \rangle)$. By the Σ_1 -elementarity of $\hat{\pi}'$, we therefore have $z = h_{\gamma, A_\nu \cap J_\gamma^X}(i, \langle \hat{\pi}'(x), \hat{\pi}'(p_\mu) \rangle)$. But $\hat{\pi}'(p_\mu) = \hat{\pi}(p_\mu) = p_\nu$ and $\hat{\pi}'(x) \in J_\lambda^X$.

In addition, $\text{sup}(H' \cap \nu) = \lambda$. That $\text{sup}(H' \cap \nu) \geq \lambda$ is clear. Conversely, let $x \in H' \cap \nu$, i.e. $x = h_{\gamma, A_\nu \cap J_\gamma^X}(i, \langle y, p_\nu \rangle)$ for some $i \in \omega$ and a $y \in J_\lambda^X$. Then x is uniquely determined by $\langle I_\gamma^0, A_\nu \cap J_\gamma^X \rangle \models (\exists z)\psi_i(z, x, \langle y, p_\nu \rangle)$. But such a z exists already in a $H_\nu^0(\alpha, \tau)$ where $H_\nu^0(\alpha, \tau)$ is the closure of S_α under all $h_{\tau, X \cap S_\tau, A_\nu \cap S_\tau}$. Since $\gamma = \text{sup}(\text{rng}(\hat{\pi}))$ and $\lambda = \text{sup}(\text{rng}(\pi))$ we can pick such $\tau \in \text{rng}(\hat{\pi})$ and $\alpha \in \text{rng}(\pi)$. Let $\bar{\tau} = \hat{\pi}^{-1}(\tau)$ and $\bar{\alpha} = \hat{\pi}^{-1}(\alpha)$. Let $\vartheta = \text{sup}(\nu \cap H_\nu^0(\alpha, \tau))$ and $\bar{\vartheta} = \text{sup}(\mu \cap H_\mu^0(\bar{\alpha}, \bar{\tau}))$. Since ν is regular in $I_{\rho(\nu)}$, $\vartheta < \nu$. Analogously, $\bar{\vartheta} < \mu$. But of course $\hat{\pi}(\bar{\vartheta}) = \vartheta$. So $x < \vartheta = \hat{\pi}(\bar{\vartheta}) < \text{sup}(\hat{\pi}[\mu]) = \lambda$.

If $\nu \in S^+$, we may define H' as $h_{\gamma, A_\nu \cap J_\gamma^X}[\omega \times (J_{\alpha_\nu}^X \times \{p_\nu\})]$ and still conclude that $H = \text{rng}(\hat{\pi}') \subseteq H'$ and $\text{sup}(H' \cap \nu) = \lambda$ by the definition of p_ν .

By lemma 19, $\hat{\pi} : \langle I_\rho^0, A \rangle \rightarrow \langle I_{\rho(\nu)}^0, A_\nu \rangle$ may be extended to a $\Sigma_{n(\nu)-1}$ -elementary embedding $\tilde{\pi}_1 : I_\beta \rightarrow I_{\beta(\nu)}$ such that ρ is the $(n(\nu)-1)$ -th projectum of I_β and A is the $(n(\nu)-1)$ -th standard code of it. Let $\hat{\pi}_0 = \hat{\pi}_1^{-1} \circ \hat{\pi}$. Then $\hat{\pi}_0 : \langle I_\rho^0, A_\mu \rangle \rightarrow \langle I_\rho^0, A \rangle$ is Σ_0 -elementary, by the coherence of $L_\kappa[X]$, and cofinal. Thus it is Σ_1 -elementary by lemma 18. Applying again lemma 19, we get a $\Sigma_{n(\nu)}$ -elementary $\tilde{\pi}_0 : I_{\beta(\mu)} \rightarrow I_\beta$.

As in lemma 24, it suffices to prove $\beta = \beta(\lambda)$, $n(\nu) = n(\lambda)$, $\rho = \rho(\lambda)$, $A = A_\lambda$, $\hat{\pi}_1^{-1}(p_\nu) = p_\lambda$, $\hat{\pi}_1^{-1}(P_\nu^*) = P_\lambda^*$, $\alpha_\nu^* = \alpha_\lambda^*$ and $\hat{\pi}_1^{-1}(\alpha_{\mu_\nu}^{**}) = \alpha_{\mu_\lambda}^{**}$. So, if $n(\nu) > 1$, we have to show that λ is $\Sigma_{n(\nu)-1}$ -regular over I_β . If $n(\nu) = 1$, then $I_\beta \models (\lambda \text{ regular})$ suffices. In addition, λ must be $\Sigma_{n(\nu)}$ -singular over I_β . For regularity, consider $\tilde{\pi}_0$ and, as in lemma 24, the least $x \in \lambda$ proving the opposite if such an x exists. This is again Σ_n -definable and therefore in $\text{rng}(\tilde{\pi}_0)$. But then $\tilde{\pi}_0^{-1}(x)$ had the same property in $I_{\beta(\mu)}$. Contradiction!

Now, assume $\nu \in S^+$. Since $I_\nu \models (\alpha_\nu \text{ is the largest cardinal})$, $H' \cap \nu$ is transitive. Thus $H' \cap \nu = \lambda$. Since $\hat{\pi}_1 : \langle I_\rho^0, A \rangle \rightarrow \langle I_\gamma^0, A \cap J_\gamma^X \rangle$ is Σ_1 -elementary and $\lambda \subseteq H' = \text{rng}(\hat{\pi}_1)$, we have $\lambda = \lambda \cap h_{\rho, A}[\omega \times (J_{\alpha_\nu}^X \times \{\hat{\pi}_1^{-1}(p_\nu)\})]$. I.e. there is a Σ_1 -map over $\langle I_\rho, A \rangle$ from α_ν onto λ . But this is then $\Sigma_{n(\nu)}$ -definable over I_β and λ is $\Sigma_{n(\nu)}$ -singular over I_β .

If $\nu \notin S^+$, then the fact that λ is $\Sigma_{n(\nu)}$ -singular over I_β , $\alpha_\nu^* = \alpha_\lambda^*$ and $\hat{\pi}_1^{-1}(\alpha_{\mu_\nu}^{**}) = \alpha_{\mu_\lambda}^{**}$ may be seen as in lemma 24 because $\pi_0(\alpha_\mu^*) = \alpha_\nu^* \in \text{rng}(\pi_0)$.

That $\hat{\pi}_1^{-1}(p_\nu) = p_\lambda$ and $\hat{\pi}_1^{-1}(P_\nu^*) = P_\lambda^*$ can again be proved as in (5) in the proof of lemma 19. \square

Lemma 26

Let $\nu \in E$ and $\Lambda(\xi, \nu) = \{\text{sup}(h_{\nu, B_\nu}[\omega \times (J_\beta^X \times \{\xi\})] \cap \nu) < \nu \mid \beta \in \text{Lim} \cap \nu\}$. Let $\bar{\eta} < \bar{\nu}$ and $\pi : \langle I_{\bar{\nu}}^0, B \rangle \rightarrow \langle I_\nu^0, B_\nu \rangle$ be Σ_1 -elementary. Then $\Lambda(\bar{\xi}, \bar{\nu}) \cap \bar{\eta} \in J_{\bar{\nu}}^X$ and $\pi(\Lambda(\bar{\xi}, \bar{\nu}) \cap \bar{\eta}) = \Lambda(\xi, \nu) \cap \pi(\bar{\eta})$ where $\pi(\bar{\xi}) = \xi$ and $\pi(\bar{\eta}) = \eta$.

Proof:

(1) Let $\lambda \in \Lambda(\xi, \nu)$. Then $\Lambda(\xi, \lambda) = \Lambda(\xi, \nu) \cap \lambda$.

Let β_0 be minimal such that

$$\sup(h_{\nu, B_{\nu}}[\omega \times (J_{\beta_0}^X \times \{\xi\})] \cap \nu) = \lambda.$$

Then, by lemma 25, for all $\beta \leq \beta_0$

$$h_{\lambda, B_{\lambda}}[\omega \times (J_{\beta}^X \times \{\xi\})] = h_{\nu, B_{\nu}}[\omega \times (J_{\beta}^X \times \{\xi\})]$$

and for all $\beta_0 \leq \beta$

$$\begin{aligned} h_{\lambda, B_{\lambda}}[\omega \times (J_{\beta_0}^X \times \{\xi\})] &\subseteq h_{\lambda, B_{\lambda}}[\omega \times (J_{\beta}^X \times \{\xi\})] \\ &\subseteq h_{\nu, B_{\nu}}[\omega \times (J_{\beta}^X \times \{\xi\})]. \end{aligned}$$

So $\Lambda(\xi, \lambda) = \Lambda(\xi, \nu) \cap \lambda$.

(2) $\Lambda(\bar{\xi}, \bar{\nu}) \cap \bar{\eta} \in J_{\bar{\nu}}^X$

Let $\bar{\lambda} := \sup(\Lambda(\bar{\xi}, \bar{\nu}) \cap \bar{\eta} + 1)$. Then, by (1), $\Lambda(\bar{\xi}, \bar{\nu}) \cap \bar{\eta} + 1 = \Lambda(\bar{\xi}, \bar{\lambda}) \cup \{\bar{\lambda}\}$. But $\Lambda(\bar{\xi}, \bar{\lambda})$ is definable over $I_{\beta(\bar{\lambda})}$. Since $\beta(\bar{\lambda}) < \bar{\nu}$, we get $\Lambda(\bar{\xi}, \bar{\nu}) \cap \bar{\eta} + 1 \in J_{\bar{\nu}}^X$.

(3) Let $\sup(h_{\bar{\nu}, B_{\bar{\nu}}}[\omega \times (J_{\bar{\beta}}^X \times \{\bar{\xi}\})]) < \bar{\nu}$ and $\pi(\bar{\beta}) = \beta$. Then

$$\pi(\sup(h_{\bar{\nu}, B_{\bar{\nu}}}[\omega \times (J_{\bar{\beta}}^X \times \{\bar{\xi}\})] \cap \bar{\nu})) = \sup(h_{\nu, B_{\nu}}[\omega \times (J_{\beta}^X \times \{\xi\})] \cap \nu).$$

Let $\bar{\lambda} := \sup(h_{\bar{\nu}, B_{\bar{\nu}}}[\omega \times (J_{\bar{\beta}}^X \times \{\bar{\xi}\})] \cap \bar{\nu})$. Then $\langle I_{\bar{\nu}}^0, B_{\bar{\nu}} \rangle \models \neg(\exists \bar{\lambda} < \theta)(\exists i \in \omega)(\exists \xi_i < \bar{\beta})(\theta = h_{\bar{\nu}, B_{\bar{\nu}}}(i, \langle \xi_i, \bar{\xi} \rangle))$. So $\langle I_{\nu}^0, B_{\nu} \rangle \models \neg(\exists \lambda < \theta)(\exists i \in \omega)(\exists \xi_i < \beta)(\theta = h_{\nu, B_{\nu}}(i, \langle \xi_i, \xi \rangle))$ where $\pi(\bar{\lambda}) = \lambda$. I.e. $\sup(h_{\nu, B_{\nu}}[\omega \times (J_{\beta}^X \times \{\xi\})] \cap \nu) \leq \lambda$. But $(\pi \upharpoonright J_{\bar{\lambda}}^X) : \langle I_{\bar{\lambda}}^0, B_{\bar{\lambda}} \rangle \rightarrow \langle I_{\lambda}^0, B_{\lambda} \rangle$ is elementary. So, if $\langle I_{\bar{\lambda}}^0, B_{\bar{\lambda}} \rangle \models (\forall \eta)(\exists \xi_i \in \bar{\beta})(\exists n \in \omega)(\eta \leq h_{\bar{\lambda}, B_{\bar{\lambda}}}(n, \langle \xi_i, \bar{\xi} \rangle))$, then $\langle I_{\lambda}^0, B_{\lambda} \rangle \models (\forall \eta)(\exists \xi_i \in \beta)(\exists n \in \omega)(\eta \leq h_{\lambda, B_{\lambda}}(n, \langle \xi_i, \xi \rangle))$. But by lemma 25, $h_{\lambda, B_{\lambda}}[\omega \times (J_{\beta}^X \times \{\xi\})] \subseteq h_{\nu, B_{\nu}}[\omega \times (J_{\beta}^X \times \{\xi\})]$. I.e. it is indeed $\lambda = \sup(h_{\nu, B_{\nu}}[\omega \times (J_{\beta}^X \times \{\xi\})] \cap \nu)$.

(4) $\pi(\Lambda(\bar{\xi}, \bar{\nu}) \cap \bar{\eta}) = \Lambda(\xi, \nu) \cap \pi(\bar{\eta})$

For $\bar{\lambda} \in \Lambda(\bar{\xi}, \bar{\nu})$,

$$\pi(\Lambda(\bar{\xi}, \bar{\nu}) \cap \bar{\lambda})$$

by (1)

$$= \pi(\Lambda(\bar{\xi}, \bar{\lambda}))$$

by Σ_1 -elementarity of π

$$= \Lambda(\xi, \pi(\bar{\lambda}))$$

by (1) and (3)

$$= \Lambda(\xi, \nu) \cap \pi(\bar{\lambda}).$$

So, if $\Lambda(\bar{\xi}, \bar{\nu})$ is cofinal in $\bar{\nu}$, then we are finished. But if there exists $\bar{\lambda} := \max(\Lambda(\bar{\xi}, \bar{\nu}))$, then, by (1) and (2), $\Lambda(\bar{\xi}, \bar{\nu}) \in J_{\bar{\nu}}^X$, and it suffices to show $\pi(\Lambda(\bar{\xi}, \bar{\nu})) = \Lambda(\xi, \nu)$. To this end, let $\bar{\beta}$ be maximal such that $\bar{\lambda} = \sup(h_{\bar{\nu}, B_{\bar{\nu}}}[\omega \times (J_{\bar{\beta}}^X \times \{\bar{\xi}\})] \cap \bar{\nu})$. I.e. $h_{\bar{\nu}, B_{\bar{\nu}}}[\omega \times (J_{\bar{\beta}+1}^X \times \{\bar{\xi}\})]$ is cofinal in $\bar{\nu}$. So, since $\pi[h_{\bar{\nu}, B_{\bar{\nu}}}[\omega \times (J_{\bar{\beta}+1}^X \times \{\bar{\xi}\})]] \subseteq h_{\nu, B_{\nu}}[\omega \times (J_{\beta+1}^X \times \{\xi\})]$ where $\pi(\bar{\beta}) = \beta$, $\sup(\text{rng}(\pi) \cap \nu) \leq \sup(h_{\nu, B_{\nu}}[\omega \times (J_{\beta+1}^X \times \{\xi\})] \cap \nu)$. Hence indeed $\pi(\Lambda(\bar{\xi}, \bar{\nu})) = \Lambda(\xi, \nu)$. \square

Lemma 27

Let $\nu \in E$, $H \prec_1 \langle I_{\nu}^0, B_{\nu} \rangle$ and $\lambda = \sup(H \cap \nu)$. Let $h : I_{\bar{\lambda}}^0 \rightarrow I_{\lambda}^0$ be Σ_1 -elementary and $H \subseteq \text{rng}(h)$. Then $\lambda \in E$ and $h : \langle I_{\bar{\lambda}}^0, B_{\bar{\lambda}} \rangle \rightarrow \langle I_{\lambda}^0, B_{\lambda} \rangle$ is Σ_1 -elementary.

Proof: By lemma 25, $B_{\lambda} = B_{\nu} \cap J_{\lambda}^X$. So it suffices, by lemma 24, to show

$rng(h) \prec_1 \langle I_\lambda^0, B_\lambda \rangle$. Let $x_i \in rng(h)$ and $\langle I_\lambda^0, B_\lambda \rangle \models (\exists z)\psi(z, x_i)$ for a Σ_0 formula ψ . Then we have to prove that there exists a $z \in rng(h)$ such that $\langle I_\lambda^0, B_\lambda \rangle \models \psi(z, x_i)$. Since $\lambda = \sup(H \cap \nu)$, there is a $\eta \in H \cap Lim$ such that $\langle I_\eta^0, B_\lambda \cap J_\eta^X \rangle \models (\exists z)\psi(z, x_i)$. And since $H \prec_1 \langle I_\nu^0, B_\nu \rangle$, we have $\langle I_\eta^0, B_\lambda \cap J_\eta^X \rangle \in H \subseteq rng(h)$. So also

$$rng(h) \models (\langle I_\eta^0, B_\lambda \cap J_\eta^X \rangle \models (\exists z)\psi(z, x_i))$$

because $rng(h) \prec_1 I_\lambda^0$. Hence there is a $z \in rng(h)$ such that $\langle I_\eta^0, B_\lambda \cap J_\eta^X \rangle \models \psi(z, x_i)$. I.e. $\langle I_\lambda^0, B_\lambda \rangle \models \psi(z, x_i)$. \square

Lemma 28

Let $f : \bar{\nu} \Rightarrow \nu$, $\bar{\nu} \sqsubset \bar{\tau} \sqsubseteq \mu_{\bar{\nu}}$ and $f(\bar{\tau}) = \tau$. If $\bar{\tau} \in S^+ \cup \widehat{S}$ is independent, then $(f \upharpoonright J_{\alpha_{\bar{\tau}}}^D) : \langle J_{\alpha_{\bar{\tau}}}^D, D_{\alpha_{\bar{\tau}}}, K_{\bar{\tau}} \rangle \rightarrow \langle J_{\alpha_\tau}^D, D_{\alpha_\tau}, K_\tau \rangle$ is Σ_1 -elementary.

Proof: If $\bar{\tau} = \mu_{\bar{\tau}} < \mu_{\bar{\nu}}$, then the claim holds since $|f| : I_{\mu_{\bar{\nu}}} \rightarrow I_{\mu_\nu}$ is Σ_1 -elementary. If $\mu_{\bar{\tau}} = \mu_\nu$ and $n(\tau) = n(\nu)$, then $P_\tau \subseteq P_\nu$. I.e. τ is dependent on ν . Thus $\bar{\tau}$ is not independent. So let $\mu := \mu_{\bar{\tau}} = \mu_\nu$, $n := n(\tau) < n(\nu)$ and $\tau \in S^+ \cup \widehat{S}$ be independent. Then, by the definition of the parameters, α_τ is the n -th projectum of μ .

Let

$$\gamma_\beta := \text{crit}(f_{(\beta, 0, \tau)}) < \alpha_\tau$$

for a β and

$$H_\beta := \text{the } \Sigma_n\text{-hull of } \beta \cup P_\tau \cup \{\alpha_\mu^*, \tau\} \text{ in } I_\mu.$$

I.e. $H_\beta = h_\mu^n[\omega \times (J_\beta^X \times \{\alpha'_\mu, \tau', P'_\tau\})]$ where

$$\alpha'_\mu := \text{minimal such that } h_\mu^n(i, \alpha'_\mu) = \alpha_\mu^* \text{ for an } i \in \omega$$

$$P'_\tau := \text{minimal such that } h_\mu^n(i, P'_\tau) = P_\tau \text{ for an } i \in \omega$$

$$\tau' := \text{minimal such that } h_\mu^n(i, \tau') = \tau \text{ for an } i \in \omega \text{ (rsp. } \tau' := 0 \text{ for } \tau = \mu).$$

For the standard parameters are in P_τ .

so H_β is Σ_n -definable over I_μ with the parameters $\{\beta, \tau, \alpha_\mu^*\} \cup P_\tau$. Let

$$\rho := \alpha_\tau = \text{the } n\text{-th projectum of } \mu$$

$$A := \text{the } n\text{-th standard code of } \mu$$

$$p := \langle \alpha'_\mu, \tau', P'_\tau \rangle.$$

So $H_\beta \cap J_\rho^X$ is Σ_0 -definable over $\langle I_\rho^0, A \rangle$ with parameters β and p . (fine structure theory!)

And γ_β is defined by

$$\gamma_\beta \notin H_\beta \quad \text{and} \quad (\forall \delta \in \gamma_\beta)(\delta \in H_\beta).$$

I.e. γ_β is also Σ_0 -definable over $\langle I_\rho^0, A \rangle$ with parameters β and p .

Let $f_0 := f_{(\beta, 0, \tau)}$ for a β , $\bar{\tau}_0 := d(f_0) < \alpha_\tau$ and $\gamma := \text{crit}(f_0) < \alpha_\tau$. Let $f_1 := f_{(\beta, \gamma, \tau)}$, $\bar{\tau}_1 := d(f_1) < \alpha_\tau$ and $\delta := \text{crit}(f_1) < \alpha_\tau$. Then $\mu_{\bar{\tau}_1}$ is the direct successor of $\mu_{\bar{\tau}_0}$ in K_τ . So $f_{(\beta, \gamma, \bar{\tau}_1)} = \text{id}_{\bar{\tau}_1}$. Hence $\mu_\eta = \mu_{\bar{\tau}_1}$ holds for the minimal $\eta \in S^+ \cup S^0$ such that $\gamma < \eta \sqsubseteq \delta$. Thus

$$\mu' \in K_\tau^+ := K_\tau - (Lim(K_\tau) \cup \{min(K_\tau)\})$$

\Leftrightarrow

$(\exists \beta, \gamma, \delta, \eta)(\gamma = \gamma_\beta \text{ and } \delta = \gamma_{(\gamma_\beta+1)})$

and $\eta \in S^+ \cup S^0$ minimal such that $\gamma < \eta \sqsubseteq \delta$ and $\mu' = \mu_\eta$)

Therefore, K_τ^+ is Σ_1 -definable over $\langle I_\rho^0, A \rangle$ with parameter p .

Now, consider $\langle I_{\alpha_\tau}^0, K_\tau \rangle \models \varphi(x)$ where φ is a Σ_1 formula. Then, since K_τ is unbounded in α_τ ,

$$\langle I_{\alpha_\tau}^0, K_\tau \rangle \models \varphi(x)$$

\Leftrightarrow

$$(\exists \gamma)(\gamma \in K_\tau^+ \text{ and } \langle I_{\alpha_\gamma}^0, K_\gamma \rangle \models \varphi(x)).$$

So $\langle I_{\alpha_\tau}^0, K_\tau \rangle \models \varphi(x)$ is Σ_1 over $\langle I_\rho^0, A \rangle$ with parameter p , resp. Σ_{n+1} over I_μ with parameters $\alpha_\mu^*, \tau, P_\tau$. But since $n = n(\tau) < n(\nu)$, f is at least Σ_{n+1} -elementary. In addition $f(\alpha_{\bar{\tau}}^*) = \alpha_\tau^*$, $f(\bar{\tau}) = \tau$, $f(P_{\bar{\tau}}) = P_\tau$. So, for $x \in \text{rng}(f)$, $\langle I_{\alpha_{\bar{\tau}}}^0, K_{\bar{\tau}} \rangle \models \varphi(f^{-1}(x))$ holds iff $\langle I_{\alpha_\tau}^0, K_\tau \rangle \models \varphi(x)$. \square

Theorem 29

$\mathfrak{M} := \langle S, \triangleleft, \mathfrak{F}, D \rangle$ is a κ -standard morass.

Proof: Set

$$\sigma_{(\xi, \nu)}(i) = h_\nu^{n(\nu)}(i, \langle \xi, \alpha_\nu^*, p_\nu \rangle).$$

Then D is uniquely determined by the axioms of standard morasses and

- (1) D^ν is uniformly definable over $\langle J_\nu^X, X \upharpoonright \nu, X_\nu \rangle$
- (2) X_ν is uniformly definable over $\langle J_\nu^D, D_\nu, D^\nu \rangle$.

(1) is clear. For (2), assume first that $\nu \in \widehat{S}$ and $f_{(0, q_\nu, \nu)} = id_\nu$. Since the set $\{i \mid \sigma_{(q_\nu, \nu)}(i) \in X_\nu\}$ is $\Sigma_{n(\nu)}$ -definable over $\langle J_\nu^X, X \upharpoonright \nu, X_\nu \rangle$ with the parameters $p_\nu, \alpha_\nu^*, q_\nu$, there is a $j \in \omega$ such that

$$\sigma_{(q_\nu, \nu)}(\langle i, j \rangle) \text{ existiert} \Leftrightarrow \sigma_{(q_\nu, \nu)}(i) \in X_\nu.$$

Using this j , we have

$$X_\nu = \{\sigma_{(q_\nu, \nu)}(i) \mid \langle i, j \rangle \in \text{dom}(\sigma_{(q_\nu, \nu)})\}.$$

So, in case that $f_{(0, q_\nu, \nu)} = id_\nu$, there is the desired definition of X_ν .

Let $\nu \in \widehat{S}$, $f_{(0, q_\nu, \nu)} : \bar{\nu} \Rightarrow \nu$ cofinal and $f(\bar{q}) = q_\nu$. Then $f_{(0, \bar{q}, \bar{\nu})} = id_{\bar{\nu}}$. And by lemma 6 (b) of [Irr2], $\bar{q} = q_{\bar{\nu}}$. So, if $\bar{\nu} = \nu$, then $f_{(0, q_\nu, \nu)} = id_\nu$. Thus let $\bar{\nu} < \nu$. Then $f_{(0, q_\nu, \nu)}(x) = y$ is defined by: There is a $\bar{\nu} \leq \nu$ such that, for all $r, s \in \omega$,

$$\sigma_{(q_{\bar{\nu}}, \bar{\nu})}(r) \leq \sigma_{(q_{\bar{\nu}}, \bar{\nu})}(s) \Leftrightarrow \sigma_{(q_\nu, \nu)}(r) \leq \sigma_{(q_\nu, \nu)}(s)$$

holds and for all $z \in J_{\bar{\nu}}^X$ there is an $s \in \omega$ such that

$$z = \sigma_{(q_{\bar{\nu}}, \bar{\nu})}(s)$$

and there is an $s \in \omega$ such that

$$\sigma_{(q_{\bar{\nu}}, \bar{\nu})}(s) = x \Leftrightarrow \sigma_{(q_\nu, \nu)}(s) = y$$

And since $\langle J_\nu^X, X_\nu \rangle$ is rudimentary closed,

$$X_\nu = \bigcup \{f(X_{\bar{\nu}} \cap \eta) \mid \eta < \bar{\nu}\}.$$

Finally, if $\nu \in \widehat{S}$ and $f_{(0, q_\nu, \nu)}$ is not cofinal in ν , then C_ν is unbounded in ν and

$$X_\nu = \bigcup \{X_\lambda \mid \lambda \in C_\nu\}$$

by the coherence of $L_\kappa[X]$.

So (2) holds. From this, (DF)⁺ follows.

By (1) and (2), $J_\nu^X = J_\nu^D$ for all $\nu \in \text{Lim}$, and for all $H \subseteq J_\nu^X = J_\nu^D$

$$H \prec_1 \langle J_\nu^X, X \upharpoonright \nu \rangle \Leftrightarrow H \prec_1 \langle J_\nu^D, D_\nu \rangle.$$

Now, we check the axioms.

(MP) and (MP)⁺

$|f_{(0, \xi, \nu)}|$ is the uncollapse of $h_{\mu_\nu}^{n[\nu]}[\omega \times \{\xi^*, \nu^*, \alpha_\nu^*, \alpha_{\mu_\nu}^{**}, P_\nu^*\}^{<\omega}]$ where ξ^* is minimal such that $h_{\mu_\nu}^{n(\nu)-1}(i, \xi^*) = \xi$. Therefore, (MP) and (MP)⁺ hold.

(LP1)

holds by (2) above.

(LP2)

This is lemma 26.

(CP1) and (CP1)⁺

This follows from lemma 24 and the definition of $\sigma_{(\xi, \nu)}$.

(CP2)

This is lemma 27.

(CP3) and (CP3)⁺

Let $x \in J_\nu^X$, $i \in \omega$ and $y = h_{\nu, B_\nu}(i, x)$. Since C_ν is unbounded in ν , there is a $\lambda \in C_\nu$ such that $x, y \in J_\lambda^X$. By lemma 25, $B_\lambda = B_\nu \cap J_\lambda^X$. So $y = h_{\lambda, B_\lambda}(i, x)$.

(DP1)

holds by the definition of μ_ν .

(DF)

Let $\mu := \mu_\nu$, $k := n(\mu)$ and

$$\pi(n, \beta, \xi) := \text{the uncollapse of } h_\mu^{k+n}[\omega \times (J_\beta^X \times \{\alpha_\mu^{**}, p_\mu^*, \xi^*\}^{<\omega})]$$

where

$$\xi^* := \text{minimal such that } h_\mu^{k+n-1}(i, \xi^*) = \xi \text{ for an } i \in \omega$$

$$p_\mu^* := \text{minimal such that } h_\mu^{k+n-1}(i, p_\mu^*) = p_\mu \text{ for some } i \in \omega$$

$$\alpha_\mu^{**} := \text{minimal such that } h_\mu^{k+n-1}(i, \alpha_\mu^{**}) = \alpha_\mu^* \text{ for some } i \in \omega.$$

Prove

$$|f_{(\beta, \xi, \mu)}^{1+n}| = \pi(n, \beta, \xi).$$

for all $n \in \omega$ by induction.

For $n = 0$, this holds by definition of $f_{(\beta, \xi, \mu)}^1 = f_{(\beta, \xi, \mu)}$. So assume that |

$f_{(\beta, \xi, \mu)}^m \mid = \pi(m-1, \beta, \xi)$ is already proved for all $1 \leq m \leq n$. Then, by definition of $\tau(m, \mu)$,

$\alpha_{\tau(m, \mu)}$ = the $(k+m-1)$ -th projectum of μ .

Let $\pi(n, \beta, \xi) : I_{\bar{\mu}} \rightarrow I_{\mu}$. Then

(*) $\xi(m, \mu) = \pi(n, \beta, \xi)\xi(m, \bar{\mu})$ for all $1 \leq m \leq n$:

Let $\pi := \pi(n, \beta, \xi)$, $\alpha := \pi^{-1}[\alpha_{\tau(m, \mu)} \cap \text{rng}(\pi)]$, $\rho := \pi(\alpha)$

$r :=$ minimal such that $h_{\mu}^{k+m-2}(i, r) = p_{\mu}$ for an $i \in \omega$

$\alpha' :=$ minimal such that $h_{\mu}^{k+m-2}(i, \alpha') = \alpha_{\mu}^*$ for an $i \in \omega$

$p :=$ the $(k+m-1)$ -th parameter of μ

and $\pi(\bar{r}) = r$, $\pi(\bar{p}) = p$, $\pi(\bar{\alpha}') = \alpha'$. Let $\bar{\xi} := \xi(m, \bar{\mu})$. Then $\bar{p} = h_{\bar{\mu}}^{k+m-1}(i, \langle \bar{x}, \bar{\xi}, \bar{r}, \bar{\alpha}' \rangle)$ for a $\bar{x} \in J_{\bar{\alpha}}^X$, because $\alpha = \alpha_{\tau(m, \bar{\mu})}$. So $p = h_{\mu}^{k+m-1}(i, \langle x, \xi, r, \alpha' \rangle)$ where $\pi(\bar{x}) = x$ and $\pi(\bar{\xi}) = \xi$. Thus $h_{\mu}^{k+m-1}[\omega \times (J_{\alpha_{\tau(m, \mu)}}^X \times \{\alpha', r, \xi\}^{<\omega})] = J_{\mu}^X$ by definition of p . So $\xi(m, \mu) \leq \xi$. Assume $\xi(m, \mu) < \xi$. Then $I_{\mu} \models (\exists \eta < \xi)(\exists i \in \omega)(\exists x \in J_{\rho}^X)(\xi = h_{\mu}^{k+m-1}(i, \langle x, \eta, r, \alpha' \rangle))$. So $I_{\bar{\mu}} \models (\exists \eta < \bar{\xi})(\exists i \in \omega)(\exists x \in J_{\bar{\alpha}}^X)(\bar{\xi} = h_{\bar{\mu}}^{k+m-1}(i, \langle x, \eta, \bar{r}, \bar{\alpha}' \rangle))$. But this contradicts the definition of $\bar{\xi} = \xi(m, \bar{\mu})$.

So, for all $1 \leq m \leq n$,

$$\xi(m, \mu) \in \text{rng}(\pi(n, \beta, \xi)).$$

In addition, for all $\beta < \alpha_{\tau(m, \mu)}$,

$$d(f_{(\beta, \xi(m, \mu), \mu)}^m) < \alpha_{\tau(m, \mu)}.$$

Consider $\pi := \pi(m-1, \beta, \xi) \mid f_{(\beta, \xi, \mu)}^m \mid$ where $\xi = \xi(m, \mu)$. Then $\pi : I_{\bar{\mu}} \rightarrow I_{\mu}$ is the uncollapse of $h_{\mu}^{k+m-1}[\omega \times (\beta \times \{\xi, \alpha', r\}^{<\omega})]$ where

$r :=$ minimal such that $h_{\mu}^{k+m-2}(i, r) = p_{\mu}$ for some $i \in \omega$

$\alpha' :=$ minimal such that $h_{\mu}^{k+m-2}(i, \alpha') = \alpha_{\mu}^*$ for some $i \in \omega$.

And $h_{\bar{\mu}}^{k+m-1}[\omega \times (\beta \times \{\bar{\xi}, \bar{\alpha}', \bar{r}\}^{<\omega})] = J_{\bar{\mu}}^X$ where $\pi(\bar{\xi}) = \xi$, $\pi(\bar{\alpha}') = \alpha'$ and $\pi(\bar{r}) = r$. Assume $\alpha_{\tau(m, \mu)} \leq \bar{\mu} < \mu$. Then there were a function over $I_{\bar{\mu}}$ from $\beta < \alpha_{\tau(m, \mu)}$ onto $\alpha_{\tau(m, \mu)}$. This contradicts the fact that $\alpha_{\tau(m, \mu)}$ is a cardinal in I_{μ} . If $\bar{\mu} = \mu$, then $f_{(\beta, \bar{\xi}, \mu)}^m = id_{\mu}$. This contradicts the minimality of $\tau(m, \mu)$.

Since $\xi(m, \mu) \in \text{rng}(\pi(n, \beta, \xi))$, we can prove

$$\text{rng}(\pi(n, \beta, \xi)) \cap J_{\alpha_{\tau(m, \mu)}}^D \prec_1 \langle J_{\alpha_{\tau(m, \mu)}}^D, D_{\alpha_{\tau(m, \mu)}}, K_{\mu}^m \rangle$$

for all $1 \leq m \leq n$ as in lemma 28.

We still must prove minimality.. Let $f \Rightarrow \mu$ and $\beta \cup \{\xi\} \subseteq \text{rng}(f)$ such that

$$\text{rng}(f) \cap J_{\alpha_{\tau(m, \mu)}}^D \prec_1 \langle J_{\alpha_{\tau(m, \mu)}}^D, D_{\alpha_{\tau(m, \mu)}}, K_{\mu}^m \rangle$$

$$\xi(m, \mu) \in \text{rng}(f)$$

holds for all $1 \leq m \leq n$. Show that f is Σ_{k+n} -elementary and that the first standard parameters including the $(k+n-1)$ -th are in $\text{rng}(f)$. That suffices because $\pi(n, \beta, \xi)$ is minimal.

Let p_{μ}^{k+m} be the $(k+m)$ -th standard parameter of μ .

Prove, by induction on $0 \leq m \leq n$,

$$\begin{aligned} & f \text{ is } \Sigma_{k+m}\text{-elementary} \\ & p_\mu^1, \dots, p_\mu^{k+m-1} \in \text{rng}(f). \end{aligned}$$

For $m = 0$, this is clear because $f \Rightarrow \mu$. So assume it to be proved for $m < n$ already. Then let $\alpha := \alpha_{\tau(m+1, \mu)}$ and $\bar{\alpha} = f^{-1}[\alpha \cap \text{rng}(f)]$. Consider $\pi := (f \upharpoonright J_{\bar{\alpha}}^D) : \langle J_{\bar{\alpha}}^D, D_{\bar{\alpha}}, \bar{K} \rangle \rightarrow \langle J_\alpha^D, D_\alpha, K_\mu^{m+1} \rangle$. Construct a Σ_{k+m+1} -elementary extension $\tilde{\pi}$ of π . To do so, set

$$f_\beta = f_{(\beta, \xi(m+1, \mu), \mu)}^{m+1}$$

$$\mu(\beta) = d(f_\beta)$$

$$H = \bigcup \{f_\beta[\text{rng}(\pi) \cap J_{\mu(\beta)}^D] \mid \beta < \alpha\}.$$

Then $H \cap J_\alpha^D = \text{rng}(\pi)$. For, $\text{rng}(\pi) \subseteq H \cap J_\alpha^D$ is clear because $f_\beta \upharpoonright J_\beta^D = id \upharpoonright J_\beta^D$. So let $y \in H \cap J_\alpha^D$. I.e. $y = f_\beta(x)$ for some $x \in \text{rng}(\pi)$ and a $\beta < \alpha$. Let $K^+ = K_\mu^{m+1} - \text{Lim}(K_\mu^{m+1})$ and $\beta(\eta) = \text{sup}\{\beta \mid f_{(\beta, \xi(m+1, \eta), \eta)}^{m+1} \neq id_\eta\}$. Then

$$\langle J_\alpha^D, D_\alpha, K_\mu^{m+1} \rangle \models (\exists y)(\exists \eta \in K^+)(y = f_{(\beta, \xi(m+1, \eta), \eta)}^{m+1}(x) \in J_{\beta(\eta)}^D).$$

Since $\text{rng}(\pi) \prec_1 \langle J_\alpha^D, D_\alpha, K_\mu^{m+1} \rangle$, $y = f_{(\beta, \xi(m+1, \eta), \eta)}^{m+1}(x) \in \text{rng}(\pi)$ if $x \in \text{rng}(\pi)$ for such an η . But since $y = f_{(\beta, \xi(m+1, \eta), \eta)}^{m+1}(x) \in J_{\beta(\eta)}^D$, we get $f_\beta(x) = f_{(\beta, \xi(m+1, \eta), \eta)}^{m+1}(x) \in \text{rng}(\pi)$.

Show $H \prec_{k+m+1} I_\mu$. Since $f_{(\beta, \xi, \mu)}^{m+1} = \pi(m, \beta, \xi)$, $\alpha_{\tau(m+1, \mu)}$ is the $(k+m)$ -th projectum of μ . Like in (*) above, we can show that the $(k+m)$ -th standard parameter p_μ^{k+m} of μ is in $\text{rng}(f_\beta)$. Now, let $I_\mu \models (\exists x)\varphi(x, y, p_\mu^1, \dots, p_\mu^{k+m})$ where φ is a Π_{k+m} formula and $y \in H \cap J_\alpha^D$. Since f_β is Σ_{k+m} -elementary, the following holds:

$$I_\mu \models (\exists x)\varphi(x, y, p_\mu^1, \dots, p_\mu^{k+m}) \Leftrightarrow (\exists \gamma \in K_\mu^{m+1})(\exists x)(I_\gamma \models \varphi(x, y, p_\gamma^1, \dots, p_\gamma^{k+m})).$$

And since $\text{rng}(\pi) \prec_1 \langle J_\alpha^D, D_\alpha, K_\mu^{m+1} \rangle$,

$$\text{rng}(\pi) \models (\exists \gamma \in K_\mu^{m+1})(\exists x)(I_\gamma \models \varphi(x, y, p_\gamma^1, \dots, p_\gamma^{k+m})).$$

Thus there is such an x in $\text{rng}(\pi)$ and therefore in H .

Let $\tilde{\pi}$ be the uncollapse of H . Then $\tilde{\pi}$ is Σ_{k+m} -elementary and, since $p_\mu^1, \dots, p_\mu^{k+m} \in \text{rng}(f_\beta)$ for all $\beta < \alpha$, we have $p_\mu^1, \dots, p_\mu^{k+m} \in \text{rng}(\pi) = H$. In addition, by the induction hypothesis, f is Σ_{k+m} -elementary and $p_\mu^1, \dots, p_\mu^{k+m-1} \in \text{rng}(f)$. Again as in (*) above, we can show that $p_\mu^{k+m} \in \text{rng}(f)$ using $\xi(m+1, \mu) \in \text{rng}(f)$. But since $\tilde{\pi}$ and f are the same on the $(k+m)$ -th projectum, we get $\tilde{\pi} = f$.

(SP) follows from $\mid f_{(\beta, \xi, \mu)}^{1+n} \mid = \pi(n, \beta, \xi)$, because for all $\nu \sqsubset \tau \sqsubseteq \mu_\nu$ such that $\tau \in S^+$ (rsp. $\tau = \nu$) the following holds:

$$p_\tau \in \text{rng}(\pi(n, \beta, \xi)) \Leftrightarrow \xi_\tau \in \text{rng}(\pi(n, \beta, \xi)).$$

This may again be shown as (*).

(DP2)

is like (*) in (DF).

(DP3)

(a) is clear.

(b) was already proved with (DF)⁺.

□

Theorem 30

Let $\langle X_\nu \mid \nu \in S^X \rangle$ be such that

- (1) $L[X] \models S^X = \{\beta(\nu) \mid \nu \text{ singular}\}$
- (2) $L[X]$ is amenable
- (3) $L[X]$ has condensation
- (4) $L[X]$ has coherence.

Then there is a sequence $C = \langle C_\nu \mid \nu \in \widehat{S} \rangle$ such that

- (1) $L[C] = L[X]$
- (2) $L[C]$ has condensation
- (3) C_ν is club in J_ν^C w.r.t. the canonical well-ordering $<_\nu$ of J_ν^C
- (4) $otp(\langle C_\nu, <_\nu \rangle) > \omega \Rightarrow C_\nu \subseteq \nu$
- (5) $\mu \in Lim(C_\nu) \Rightarrow C_\mu = C_\nu \cap \mu$,
- (6) $otp(C_\nu) < \nu$.

Proof: First, construct from $L[X]$ a standard morass as in theorem 29. Then construct a inner model $L[C]$ from it as in [Irr2]. □

References

- [BJW] A. Beller, R.-B. Jensen, P. Welch: **Coding the Universe**, London Mathematical Society Lecture Notes Series, Cambridge University Press, London, 1982
- [Dev1] K. Devlin: **Aspects of Constructibility**, Lecture Notes in Mathematics, no. 354, Springer-Verlag, Berlin, 1973
- [Dev2] K. Devlin: **Constructibility**, Springer-Verlag, Berlin, 1984
- [DJS] H.-D. Donder, R.-B. Jensen, L.J. Stanley: Condensation-Coherent Global Square Systems, **Recursion theory**, Proceedings of Symposia in Pure Mathematics, vol. 42, American Mathematical Society, Providence, 1985, 237–258
- [Irr1] B. Irrgang: **Kondensation und Morasse**, Dissertation, München, 2002
- [Irr2] B. Irrgang: **Proposing (ω_1, β) -Morasses**, $\omega_1 \leq \beta$
- [Jen1] R.-B. Jensen: The Fine Structure of the Constructible Hierarchy, **Annals of Mathematical Logic**, vol. 4 (1972), 229–308
- [Jen2] R.-B. Jensen: **Higher-Gap Morasses**, hand-written notes, 1972/73